

02-03-00

A

EXPRESS MAIL LABEL NO. 491077878US

Date of Deposit: February 1, 2000

WDN.AAR ck 2/1/00 E-209-99/0

PATENT

Attorney's Ref. No. 4239-53335

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Box PATENT APPLICATION
TO THE ASSISTANT COMMISSIONER FOR PATENTS
Washington, D.C. 20231

Transmitted herewith for filing is the patent application of:

Inventor(s): Dwight W. Miller, Richard Beger, Jackson O. Lay, Jr.,
Jon G. Wilkes and James P. Freeman

For: METHODS FOR PREDICTING THE BIOLOGICAL,
CHEMICAL, AND PHYSICAL PROPERTIES OF
MOLECULES FROM THEIR SPECTRAL PROPERTIES

Enclosed are:

- ☒ 69 pages of specification, 8 pages of claims, an abstract and a Combined Declaration and Power of Attorney.
- ☒ 16 sheet(s) of informal drawings.


| FILING FEE | | | | | |
|------------------------------|-----------------|----------------|-----------------|----------|-----------|
| For | Claims Filed | Number Free | Number Extra | Rate | Basic Fee |
| Total Claims | 54 | 20 | = 34 | \$18.00 | \$ 612.00 |
| Independent Claims | 7 | 3 | = 4 | \$78.00 | \$ 312.00 |
| Multiple Dependent Claim Fee | | | | \$260.00 | |
| TOTAL FILING FEE | | | | | \$1614.00 |

- ☒ A check in the amount of \$1,614.00 to cover ☒ filing fee is enclosed.
- ☒ The Commissioner is hereby authorized to charge any additional fees which may be required in connection with the filing of this application and recording any assignment filed herewith, or credit over-payment, to Account No. 02-4550. A copy of this sheet is enclosed.
- ☒ Please return the enclosed postcard to confirm that the items listed above have been received.

Respectfully submitted,

KLARQUIST SPARKMAN CAMPBELL
LEIGH & WHINSTON, LLP

By



William D. Noonan, M.D.

Registration No. 30,878

One World Trade Center, Suite 1600
121 S.W. Salmon Street
Portland, Oregon 97204
Telephone: (503) 226-7391
Facsimile: (503) 228-9446
cc: Docketing Secretary
Aden A. Rehms, Ph.D.

S:\WPDOCS\WDN\NIH\53335\Patent App Trans doc

**METHODS FOR PREDICTING THE BIOLOGICAL,
CHEMICAL, AND PHYSICAL PROPERTIES OF
MOLECULES FROM THEIR SPECTRAL PROPERTIES**

5

FIELD OF THE INVENTION

The invention relates to methods for predicting the biological, chemical, and physical properties of molecules from their spectral properties.

10

BACKGROUND

The number of known chemical compounds is vast, and the number is increasing constantly because methods for isolating and synthesizing molecules continue to improve. For instance, chemists are now able to employ the techniques of combinatorial chemistry to synthesize thousands of different chemical compounds at one time using a mixture of only a few interchangeable chemical building blocks. Furthermore, chemists are now able to use combinatorial computer models to generate large numbers of chemical structures that may in theory be synthesized.

While there are many chemical compounds, only a relative few of those compounds may exhibit a particular desirable property, such as pharmaceutical activity. Random testing of known compounds to find those that show pharmaceutical activity is very expensive and time-consuming. Similarly, there is a need to screen compounds for toxicity, so that rational decisions can be made regarding the use and regulation of compounds that have toxic potential. At present, only a fraction of known compounds have been thoroughly tested for their toxicological and potential therapeutic properties. To address this problem, scientists have developed methods which attempt to predict which compounds are likely to exhibit a particular property.

Methods for predicting the properties of chemical compounds are generally based upon the related observations that the structure of a compound is related to its biological, chemical, and physical properties, and that compounds of similar structure exhibit similar properties. These observations have been used to search for new compounds exhibiting a particular property. For instance, a benzene ring

is present in both acetaminophen and salicylamide, both of which are analgesics. Although incorporating a benzene ring into a new molecule increases the likelihood that it too will exhibit analgesic activity, this deduction only narrows the compounds to be tested. This approach is still basically one of trial and error, because many compounds with a benzene ring are not analgesics. Moreover, analgesics without a benzene ring will be missed in the search.

Quantitative structure-property relationships and quantitative structure-activity relationships (collectively QSAR) are attempts to quantify the observed relationships between the structure of chemical compounds and the extent to which those compounds exhibit certain properties. For instance, a QSAR might attempt to quantify how the analgesic activity of known analgesics that contain a benzene ring (such as acetaminophen and salicylamide) depends upon the number and identity of substituents on their benzene rings. Once established, such a QSAR could be used to predict the analgesic activity of other compounds that contain benzene rings, and identify those compounds that warrant further investigation as analgesics based on their predicted analgesic activity.

In the terminology associated with the QSAR method, the property for which a prediction is sought, such as analgesic activity, is termed the "endpoint." In general the endpoint may be any measurable biological, chemical or physical property.

To establish a QSAR, endpoint values are obtained for a set of compounds and a correlation is then sought between the endpoint values and some measure(s) of structure available for each of the compounds. The measures used to describe or reflect the structure of the compounds for which a correlation is sought are termed structure descriptors. Structure descriptors may be defined directly with reference to the known structures of the compounds, or may indirectly reflect the structure through a property of the molecule that is sensitive to changes in structure. For example, an investigator might try to correlate the analgesic activity of compounds that contain a benzene ring with either a direct measure of the structure, such as the number of hydroxyl groups attached to the benzene ring, or an indirect measure of the structure of the compounds, like water solubility. If the

direct measure is chosen, the attempted correlation could only include those compounds with hydroxyl groups on the benzene ring, while the indirect measure is more general and could be used to include all benzene ring containing compounds in the attempted correlation.

5 The endpoint data and the structure descriptor(s) for the set of compounds that are chosen to establish a QSAR are termed the training set. In general, the reliability of a QSAR increases as the number of compounds in the training set increases. If possible, the training set desirably includes compounds that exhibit a wide range of endpoint values and possess diverse structures.

10 If the endpoint and structure descriptor(s) are sufficiently correlated, a mathematical or graphical representation may be obtained. For example, the growth inhibition of certain gram negative bacteria by aromatic amines is correlated in a linear fashion with the logarithm of the octanol-water partition coefficient, and the correlation indicates that as the aromatic amines become more
15 hydrophobic they are more likely to inhibit the growth of these bacteria (Hansch and Leo, *Exploring QSAR: Fundamentals and Applications in Chemistry and Biology*, American Chemical Society, 1995, p. 416).

 The QSAR representation may be used to predict endpoint values for other compounds from their structure descriptor(s), and the reliability of a QSAR may
20 be tested using a validation set of data. A validation set includes structural and endpoint data for compounds that were not part of the training set. The validation set desirably exhibits a diversity of endpoint values and structures that is commensurate with the training set. The QSAR is tested by how reliably it predicts endpoint data for the validation set compounds from the validation set
25 structural data. For example, α -naphthylamine, an aromatic amine, is about two times less toxic than predicted by its octanol-water partition coefficient, indicating that the simple linear relationship described above is not always reliable and that some specific interaction is responsible for its behavior.

 The Hammett equation is another example of a simple QSAR that relates a
30 single structure descriptor (in this case, derived directly from the structure of the compounds) to an endpoint (in this case, the equilibrium or rate constant for a

particular type of reaction). The Hammet equation is a linear equation of the form:

$$\ln K_x = \rho\sigma + \ln K_H$$

- 5 The electronic parameter (σ) is a measure of the ability of a group of atoms (typically a substituent at a particular structural position) to donate (negative value) or withdraw (positive value) electron density to or from the reaction center of the molecule to which it is attached. The slope parameter (ρ) is a measure of the sensitivity of the reaction to the withdrawal or release of electron density, and is
- 10 constant for a particular type of reaction. K_x and K_H are, respectively, the equilibrium constant for reaction of the substituted molecule and the equilibrium constant for reaction of the unsubstituted parent molecule. The reaction rate constants, k_x and k_H , may replace the equilibrium constants. A plot of $\ln K_x$ (or $\ln k_x$) versus σ is often linear and may be used, for example, to predict the
- 15 equilibrium constant (or rate constant) for other structurally similar compounds from only the σ values of their substituents.

- The Hammet parameter σ is an example of a structure descriptor that is derived from experimental data for compounds of known structure. As described above, σ is a measure of the electronic properties of a substituent. Specifically σ
- 20 measures the ability of a substituent at a particular position to donate or withdraw electron density and may be defined according to the following equation:

$$\sigma = \log K_x - \log K_H$$

- 25 where K_x and K_H are, respectively, the acid ionization constants for an aromatically substituted benzoic acid derivative and for unsubstituted benzoic acid. The parameter σ for amino group substitution in a position para to the reaction center is determined, for example, from the measured acid ionization constants of *p*-aminobenzoic acid and benzoic acid using the above equation. Although
- 30 depending on the position of substitution, electron-withdrawing substituents generally tend to stabilize the anion formed when benzoic acid ionizes, making K_x larger than K_H and σ positive. Conversely, electron-donating substituents tend to

destabilize the anion and generally have negative σ values that are also dependent upon the position of substitution. Acid dissociation data for benzoic acid derivatives with various substituents in ortho, meta or para positions have been used to generate σ values for many substituents in these structural positions.

- 5 Unfortunately, the σ values derived for substituents in this manner are typically not valid for multiply substituted molecules because σ values may not be additive.

Due to the inability of σ values to accurately reflect the cumulative electronic effect of multiple substituents, alternative structure descriptors have been investigated. For example, Ijzerman et al. derived structure descriptors from the assigned ^{13}C NMR shifts of the aromatic carbons in a family of N-*tert*-butylphenylethanolamines (Ijzerman et al., *J. Med. Chem.*, **29**: 549-554, 1986). Ijzerman et al. subtracted the value of the chemical shift for the carbon atoms in benzene (128.5ppm) from the assigned chemical shift values for each of the carbons in the benzene ring of each of the substituted N-*tert*-butylphenylethanolamines to yield sets of structure descriptors for the compounds. A correlation was found between certain linear combinations of these structure descriptors and the β_2 -adrenoreceptor intrinsic activity of the substituted N-*tert*-butylphenylethanolamines. Since these structure descriptors are defined in a manner similar to the Hammett σ parameter, they necessarily rely upon knowledge of structure beforehand. Also, the scope of usefulness for this type of structure descriptor is limited to the compounds containing a benzene ring. Furthermore, these structure descriptors ignore the aliphatic carbon atoms of the molecules, and their contribution to the intrinsic activity (a valid assumption only when comparing the intrinsic activities of compounds that differ minimally in structure).

25 Additionally, the correlations of Ijzerman et al. have been questioned because the structure descriptors they derived from the chemical shifts of the aromatic carbons in these compounds are themselves highly correlated, and their use leads to the unrealistic result that substitution at one ortho position of the benzene ring affects intrinsic activity in a manner opposite to substitution at the other ortho position (see, Cramer et al., *Quant. Struct.-Act. Relat.*, **7**: 18-25, 1988).

30

A similar attempt to utilize spectroscopically derived parameters as structure descriptors was made by Nishikawa and Tori (Nishikawa and Tori, *J. Med. Chem.*, **27**: 1657-1663, 1984). The reactivity of the β -lactam ring of 3-substituted and 3-methylene substituted cephalosporins toward alkaline hydrolysis showed a correlation with changes in the ^{13}C NMR chemical shift of selected carbon atoms, and with changes in the infrared (IR) stretching frequency observed for the β -lactam carbon-oxygen bond. Selection of these measures was guided by knowledge of the structure of these compounds, and the role β -lactam reactivity plays in their antibiotic activity. Like the method of Ijzerman et al., the method of Nishikawa defines its structure descriptors in terms based upon a reference compound of known structure and utilizes spectral features assigned to particular atoms of known structures. Furthermore, the method, through its definition of the structure descriptors, is similarly restricted to narrow classes of compounds that differ only by simple substitutions.

Structure descriptors may be obtained theoretically or experimentally. Dipole moments and lowest unoccupied molecular orbital (LUMO) energies are examples of structure descriptors that may be obtained from theoretical quantum mechanical calculations on known structures. Experimental data correlated with a specific structural feature, common to a set of closely related compounds, may also be used to generate measures of structure (for example, the Hammett σ parameter). Bulk experimental measures of structure, such as partition coefficients (as a measure of polarity) and molar refractivities (as a measure of steric size) can also be utilized as structure descriptors. Structure descriptors based upon bulk physical properties have the advantage that they do not require structural knowledge beforehand, however such descriptors lack specificity. For example, compounds with vastly different biological activities may have very similar partition coefficients.

A particularly important type of biological QSAR uses, as an endpoint, the ability of one molecule to bind to another molecule. An example of such an endpoint would be the ability of a series of molecules to act as ligands for a regulatory protein, such as a hormone receptor. In recognition of the important

role played by three-dimensional structure, especially for biochemical reactions where molecular recognition is an important factor, the field of 3D-QSAR has emerged. The 3D-QSAR technique is exemplified by the Comparative Molecular Field Analysis (CoMFA) method of Cramer and Wold (U.S. Patent No.

5 5,025,388). The CoMFA method attempts to correlate the three-dimensional steric and electrostatic properties of a series of molecules with their relative endpoint values. The steric and electrostatic properties of a molecule are obtained from quantum mechanical or electrostatic calculations based upon known molecular structures and serve as structure descriptors. The calculations, in effect,
10 map the electron density distribution around a molecule to create a 3-D picture of its steric and electrostatic fields (collectively, the molecular field). Those steric features (e.g., bulky substituents) and/or electrostatic properties (e.g., a strong molecular dipole) that are most important in determining endpoint values are revealed by comparing the molecular fields of the molecules in the training set to
15 their endpoints. The advantage of calculated 3D-QSAR molecular field structure descriptors is that unlike structure descriptors referenced to a certain structural feature, molecular field structure descriptors enable the identification of structurally dissimilar molecules that have similar steric and electrostatic properties.

20 However, a particular problem associated with the CoMFA method and other 3D-QSAR techniques is that these methods generally require some assumptions about how molecules orient themselves relative to each other upon binding. Selecting the common alignment of a training set containing diverse structures may be problematic, leading to incorrect predictions of binding ability.
25 Furthermore, QSAR based upon quantum mechanical or electrostatic potential calculations also suffers to some extent from the inaccuracy of the calculations themselves. These calculations are by nature approximate, and become less and less reliable as molecular size increases. In addition, the effects of solvation on the quantum mechanical properties of a molecule are often difficult to calculate and
30 typically are ignored. Finally, these calculations are time-consuming and require knowledge of the molecular structures beforehand.

Isolation and structure elucidation of molecules is an expensive and time-consuming process that is desirably avoided, especially when screening large numbers of molecules (e.g., those generated in a combinatorial library).

Furthermore, many molecular structures are proprietary, so there may be
5 reluctance on the part of investigators to share training set structural and endpoint data with competitors in a common effort to improve the reliability of predictive QSARs.

It would be advantageous to have QSAR methods that utilize structure descriptors which may be obtained without knowledge of or disclosure of
10 molecular structures. Such structure descriptors could be more specific than bulk physical properties, and enable the QSAR methods to differentiate dissimilar molecular structures that exhibit similar bulk physical properties. Furthermore, the structure descriptors could avoid definition with reference to a particular structural element (e.g., the Hammett σ parameter), and therefore be useful for
15 establishing QSARs for more structurally diverse sets of compounds. Additionally, such QSAR methods could eliminate the costly step of structure elucidation and obviate the need for isolation of the subject compounds.

QSAR methods that utilize structure descriptors that are inherently reflective of the steric and electrostatic properties of molecules and the effects of
20 solvation thereon are also needed. Such methods would also desirably eliminate the necessity for assumptions regarding molecular orientation in relationship to intermolecular binding and obviate the need to rely upon approximate calculations.

SUMMARY OF THE DISCLOSURE

The present invention avoids some of the foregoing problems by providing
25 a method for predicting a biological activity of a molecule, by obtaining spectral data (such as NMR data) for a test compound, and comparing the spectral data for a test compound to a pattern derived not exclusively from the assigned spectral data (such as NMR data) of a training set of compounds having known biological activity. Similarities between the pattern of spectral data associated with the
30 biological activity of the training set compounds and the spectral data for the test compound are detected to determine whether the test compound is predicted to

exhibit the biological activity. The spectral data of the compound for which a prediction is sought need not first be correlated with corresponding structural features. Furthermore, the pattern of spectral data associated with the biological activity may be derived without first correlating the spectral data with

5 corresponding structural features. Training set patterns and similarities between the training set patterns and the test compound's spectral data are conveniently detected, in some embodiments, by segmenting the spectral data of the training set and test compounds into sub-spectral units (bins). These sub-spectral units, or bins, may be of a width corresponding to the digital resolution of the method used

10 to generate the spectral data or greater.

For example, the biological activity of a test compound may be predicted by comparing the signals in bins of the training set spectral data that are found to be associated with a biological activity (such as strong estrogen receptor binding) to signals in corresponding bins of the spectral data of the test compound.

15 Numerous signals in the test compound's spectral data that fall within bins corresponding to strong estrogen receptor binding are an indication that the test compound possesses strong estrogen receptor binding. Some of the bins of the training set spectral data may contain signals more consistently associated with strong estrogen receptor binding, in which case the presence of signals in the

20 corresponding bins for the test compound would be more heavily weighted in assigning a predicted biological activity to the test compound. Lesser numbers of signals in the test compound's spectral data that fall within bins corresponding to strong estrogen receptor binding indicate that the test compound possess only moderate or weak estrogen receptor binding.

25 The spectral data of the training compounds and the test compound may be just one type of spectral data (such as NMR, for example ^{13}C -NMR), or more than one type of spectral data (such as a composite of two or more of NMR, mass spectral, infrared, ultraviolet-visible, fluorescence, or phosphorescence data). In particular embodiments, the spectral data is a composite of two or more of nuclear

30 magnetic resonance spectroscopic (NMR) data, mass spectroscopic (MS) data,

infrared (IR) spectroscopic data, and ultraviolet-visible (UV-Vis) spectroscopic data.

In other embodiments, the spectral data of the training set compounds is segmented into sub-spectral units (bins), and scaled to normalize the importance of different signals (e.g., those from different types of spectra in a composite or those arising from signals of different inherent intensities) prior to pattern recognition. In particular embodiments, the scaling is auto-scaling. In yet other embodiments, the spectral data of the training set compounds is weighted prior to pattern recognition to emphasize those sub-spectral units (bins) that are most important for differentiating endpoint classes (such as strong versus weak estrogen receptor binding) of compounds in the training set. In more particular embodiments, the weighting is Fisher-weighting.

A pattern of spectral data associated with a biological activity can be advantageously derived from the training set spectral data using computer implemented pattern recognition techniques. Furthermore detection of similarities between the pattern derived from the training set spectral data and the spectral data exhibited by a test compound is also advantageously performed using computer implemented methods. In one such approach, the pattern of spectral data associated with a biological activity is derived for a training set of compounds by segmenting the spectral data and generating a set of canonical variate factors, one for each bin of the segmented spectral data. These canonical variate factors are used with the spectral data of a test compound to yield a prediction of the biological activity of the test compounds.

The methods of the present invention are advantageously computer implemented. In one such embodiment, the method is a computer implemented system for predicting biological activity of a test compound, in which input spectral data is received for a test compound, and for a set of training compounds having a known biological activity. The spectral pattern derived from the training set (derived, for example, using computer implemented pattern recognition programs) and the spectral pattern of the test compound are compared to determine whether the spectral patterns of the test compound match spectral patterns of the

training set associated with a biological activity. The spectral data for the test compound and the spectral data for the training set may conveniently be divided into substantially identical spectral bins, so that a signal within individual corresponding spectral bins is compared between the pattern derived from the training set and the test compound.

In this computer implemented system, the spectral patterns are obtained by inputting spectral data such as one or more of nuclear magnetic resonance data, mass spectral data, infrared data, ultraviolet-visible data, fluorescence data, and phosphorescence data. In particular embodiments, the spectral data is a composite of nuclear magnetic resonance data and mass spectral data. The spectral data of the training set is converted into principal components (PCs) and canonical variates (CVs). Peaks in particular bins of the canonical variates that are associated with a biological activity (such as high affinity to a hormone receptor) are identified, and the test compound is analyzed for the presence of one or more (and ideally many) corresponding peaks in its composite spectrum.

Another aspect of the invention is a method for predicting a biological, chemical, or physical property of molecules, by providing spectral data segmented into spectral sub-units, for a plurality of training compounds; inputting the segmented spectral data and endpoint data into a pattern-recognition program; training the pattern-recognition program with the segmented spectral data and endpoint data to establish a relationship between the spectral sub-units of the segmented spectral data and the endpoint; providing segmented spectral data for a test compound that is segmented into substantially the same spectral sub-units that were used for the training set, and comparing the relationship between the spectral sub-units of the segmented spectral data and the endpoint to the spectral sub-units of the test compound's segmented data to predict the endpoint of the test compound. In this method, the structures of the training compounds and the test compound are not necessarily known beforehand. In particular embodiments of the method, the segmented spectral data of the training set is auto-scaled and Fisher-weighted before training the pattern recognition program.

The spectral data is chosen from the group consisting of nuclear magnetic resonance data, mass spectral data, infrared data, UV-Vis data, fluorescence data, phosphorescence data, and composites thereof, for example ^{13}C NMR data, EI MS data, and composites thereof. The endpoint may be a ligand-target molecule-binding affinity, such as estrogen-receptor binding affinity. Other examples of the endpoint are a measure of biodegradability; a measure of toxicity; participation in a metabolic pathway; a partition coefficient; a reaction rate; a quantum yield, a measure of phototoxicity, an equilibrium constant; and a site of reaction on a molecular structure. In a particular embodiment, the endpoint is the octanol/water partition coefficient.

Additional non-spectral structure descriptors may be utilized along with segmented spectral data to provide an expanded set of structure descriptors useful for establishing a predictive relationship for the endpoint. Examples of non-spectral structure descriptors that do not necessarily require structural knowledge beforehand include partition coefficients, solubilities, relative acidities, relative basicities, pKa, pKb, reaction rates, and equilibrium constants. In a particular embodiment, the partition coefficient is the octanol/water partition coefficient. Calculated non-spectral descriptors are one example of descriptors requiring structural knowledge beforehand that may be utilized along with segmented spectral data in establishing a predictive relationship for an endpoint.

Another aspect of the invention is a method of using spectral data as a set of structure descriptors for a compound that does not necessarily require knowledge of the compound's structure beforehand, by providing spectral data of a training set of compounds and segmenting the spectral data into bins.

Yet another aspect of the invention is a method for establishing a relationship between spectral data and a biological, chemical, or physical property, by providing spectral data for a training set of compounds, segmenting the spectral data into bins, and detecting patterns in the bins of the spectral data that are associated with the property. The method may also include detecting corresponding patterns in spectral data of test compounds to select the test compounds having the property. Furthermore, the test compounds may be

mixtures of compounds. The spectral data of the training set can be auto-scaled and weighted (for example by Fisher-weighting) to help identify data that are most strongly associated with the biological activity. Knowledge of the structural features that lead to the spectral data is not needed beforehand.

5 Yet another aspect of the invention is a method of determining the structural features of a plurality of compounds that contribute to determining a particular endpoint property exhibited by the compounds, by providing segmented spectral data for the plurality of compounds; providing endpoint data for the plurality of compounds; establishing a spectral data-activity relationship (SDAR)
10 by identifying the segmented spectral features that bias toward increased endpoint values and the segmented spectral features that bias toward decreased endpoint values; and identifying the structural feature leading to the segmented spectral features that bias toward increased or decreased endpoint values for the plurality of compounds. As with the earlier described embodiments of the invention, the
15 segmented spectral data may be a composite of several types of spectral data.

Any of the methods of the present invention can be performed without reference to a chemical structure of the test compound. Hence the spectral features of the training set and the test compound may be compared, without determining the chemical structure of the compounds of either the training set or
20 the test compound or compounds. Any of the foregoing methods can also be incorporated into a computer readable medium, having stored thereon instructions for performing the steps of these methods.

The foregoing and other objects, features, and advantages of the invention will become more apparent from the following drawings and the detailed
25 description of several embodiments.

BRIEF DESCRIPTION OF THE DRAWINGS

Figs 1(a), 1(b) and 1(c) are the ¹³C NMR, EI MS, and IR spectra, respectively, for bisphenol A (4,4'-isopropylidenediphenol, structure shown in Fig.
30 1(a)).

Figs. 2(a), 2(b) and 2(c) show tables of data that correspond respectively to each of the spectra shown in Figs. 1(a), 1(b) and 1(c).

Fig. 3 shows a hypothetical set of structure descriptors derived from the spectral data summarized in the tables of Figs. 2(a), 2(b) and 2(c).

5 Fig. 4 shows a flowchart of a particular embodiment of the spectral data-activity relationship (SDAR) method.

Fig. 5 shows the discriminant function using ^{13}C Nuclear Magnetic Resonance (NMR) spectral data for 30 compounds in the Spectral Data-Activity Relationship (SDAR) model. The X-axis is the first canonical variate (discriminant function) and the Y-axis is the component frequency. The numbers in each box correspond to the numbers in Table 1 that serve to identify the compounds used in the SDAR. White boxes correspond to strong estrogen receptor binding compounds, and gray boxes correspond to medium estrogen receptor binding compounds.

15 Fig. 6 presents the first canonical variate factor weights using ^{13}C NMR spectral data for 30 compounds in the SDAR model. The X-axis is the bin number and the Y-axis is the factor weight relative intensity. The bins are numbered from 550 (corresponding to 0 ppm) to 770 (corresponding to 220 ppm).

Fig. 7 shows the discriminant function using composite ^{13}C NMR spectral data and Electron Impact (EI) mass spectral data for 30 compounds in the SDAR model. The X-axis is the first canonical variate and the Y-axis is the component frequency. The numbers in each box correspond to the numbers in Table 1 that serve to identify the compounds used in the SDAR. White boxes correspond to strong estrogen receptor binding compounds, and gray boxes correspond to medium estrogen receptor binding compounds.

25 Figs. 8(a) - 8(b) present the first canonical variate factor weights using composite ^{13}C NMR spectral data and EI mass spectral data for 30 compounds in the SDAR model. The X-axis is the bin number and the Y-axis is the factor weight relative intensity. EI mass spectral data are in the bins numbered from m/z 0 to 550. ^{13}C NMR spectral data occupies the bins numbered from 550 (corresponding to 0 ppm) to 770 (corresponding to 220 ppm).

Fig. 9 shows the discriminant function using ^{13}C Nuclear Magnetic Resonance (NMR) spectral data for 119 compounds in the Spectral Data-Activity Relationship (SDAR) model. The X-axis is the first canonical variate (discriminant function) and the Y-axis is the second canonical variate. The symbol S represents a strong estrogen receptor binder, M represents a medium estrogen receptor binder, and W represents a weak estrogen receptor binder.

Fig. 10 presents the first canonical variate factor weights using ^{13}C NMR spectral data for 119 compounds in the SDAR model. The X-axis is the bin number and the Y-axis is the factor weight relative intensity. The bins are numbered from 550 (corresponding to 0 ppm) to 770 (corresponding to 220 ppm).

Fig. 11 shows the discriminant function using composite ^{13}C NMR spectral data and Electron Impact (EI) mass spectral data for 119 compounds in the SDAR model. The X-axis is the first canonical variate and the Y-axis is the second canonical variate. The symbol S represents a strong estrogen receptor binder, M represents a medium estrogen receptor binder, and W represents a weak estrogen receptor binder.

Figs. 12(a) - 12(b) present the first canonical variate factor weights using composite ^{13}C NMR spectral data and EI mass spectral data for 119 compounds in the SDAR model. The X-axis is the bin number and the Y-axis is the factor weight relative intensity. EI mass spectral data are in the bins numbered from m/z 0 to 550. ^{13}C NMR spectral data occupies the bins numbered from 550 (corresponding to 0 ppm) to 770 (corresponding to 220 ppm).

Fig. 13 is a diagram of a distributed computing environment in which the present invention can be implemented.

Fig. 14 is a block diagram of a computer system that can be used to implement the present invention.

DETAILED DESCRIPTION OF SEVERAL DISCLOSED EMBODIMENTS

The illustrative embodiments that follow are best understood by first understanding terms in the art that are relevant. These terms are defined below.

Definitions and Abbreviations

LOO – leave-one-out cross validation

RBA – relative binding affinity

NMR – nuclear magnetic resonance

IR- infrared (spectroscopy)

MS-mass spectrometry

EI MS –electron impact mass spectrometry

UV-Vis –ultraviolet-visible (spectroscopy)

CODESSA – comprehensive descriptors for structural and statistical analysis

SDAR –spectral data-activity relationship

Endpoint – a particular biological, chemical, or physical property or a set of such properties for a compound that are either qualitatively or quantitatively measurable.

Structure Descriptors – any direct or indirect measure of the structure of a compound that may be obtained by theoretical or experimental means.

Training Set – endpoint data and structure descriptors (structural data) for a group of compounds used to establish a correlation between the endpoint property and the structures of the compounds.

Validation Set - endpoint data and structure descriptors (structural data) for a group of compounds used to test the reliability of the correlation between the endpoint property and the structures of the compounds.

Segmented Spectral Data – spectral data that is divided into discrete sub-spectral units (bins), each of which spans a particular spectral range. The spectral range spanned by a particular bin corresponds to a range of frequencies or a range of wavelengths for spectroscopic data and may be equal to the digital resolution of the spectral data or greater. For mass spectrometric techniques, the spectral range

within each bin corresponds to a particular mass or range of masses and may be equal to the digital resolution of the spectral data or greater. The bins need not all be of equal width. The spectral data that is divided into bins may either encompass all the spectral data of a particular type that are available or cover only a portion of the spectral data of a particular type that are available. Each bin contains information derived from the spectral signals (or lack thereof) that appear within the spectral range defined by a particular bin. The structural aspect(s) of the compounds that give rise to the information falling within any particular bin need not be known.

10 **Spectral Data-Activity Relationship (SDAR)** – a correlation between endpoint data and spectral data for a group of compounds, useful for among other things for predicting the endpoint data for compounds from their spectral data.

Nuclear Magnetic Resonance (NMR) – a phenomenon exhibited by a large number of atomic nuclei in which nuclei in a magnetic field absorb energy from a radio-frequency field at certain characteristic frequencies. Particular examples of nuclei that exhibit this phenomenon include ^{13}C , ^1H and ^{31}P .

Mass Spectrometry (MS) – a method of chemical analysis in which the substance to be analyzed is placed in a vacuum and reduced to low pressure. The resulting vapor is exposed, for example, to a beam of electrons which causes ionization to occur, either of the molecules or their fragments. The ions thus produced are accelerated and then passed through a mass analyzer that separates the ions according to their mass.

Electron Impact Mass Spectrometry (EI MS) – a mass spectrometric technique in which the ionization of molecules and their fragments is accomplished by a beam of electrons that impacts the molecules and their fragments. Typically, as the energy of the electron beam is increased, the number of fragments produced from a molecule increases.

Infrared Spectroscopy (IR) – an analytical technique which measures a range of wavelengths (or frequencies) in the infrared region or near-infrared region of the electromagnetic spectrum that are absorbed by a specimen, which characterize its molecular constitution. Infrared absorption bands identify

molecular structure components, such as aromatic, olefin, aliphatic, aldehyde, ketone, carboxylic acid, alcohol, amine, and amide groups. The frequency at which absorption occurs also reflects the frequency at which the bonds in these components stretch and and/or bend.

5 **Ultraviolet-Visible Spectroscopy (UV-Vis)** – an analytical technique which measures a range of wavelengths (or frequencies) in the ultraviolet and visible regions of the electromagnetic spectrum that are absorbed by a specimen, which characterize the electronic energy levels of its molecular constituents. UV-Vis absorption bands may be characteristic of certain molecular components, such as
10 aromatic groups or carboxyl (CO) groups.

Fluorescence Spectroscopy – an analytical technique which measures a range of wavelengths (or frequencies) of light a molecule emits in passing from a higher to lower energy electronic state during a given time period (such as the first
15 millisecond) after absorbing a photon of light. Fluorescence wavelengths and emission intensity reflect the redistribution of energy in the molecule after light absorption. Fluorescence excitation spectroscopy reflects the efficiency with which a molecule converts absorbed energy into fluorescent emission as a function of the wavelength of the absorbed photons.

Phosphorescence Spectroscopy - an analytical technique which measures a
20 range of wavelengths (or frequencies) of light a molecule emits in passing from a higher to lower energy electronic state on a time scale beyond the first millisecond after absorbing a photon of light. Phosphorescence wavelengths and emission intensity also reflect the redistribution of energy in the molecule after light absorption. Phosphorescence excitation spectra reflect the efficiency with which a
25 molecule converts absorbed energy into phosphorescent emission as a function of the wavelength of the absorbed photons.

Principal Component Analysis (PCA) – this pattern recognition technique and the related technique of partial least squares analysis reduce the structure
30 descriptors used in SDAR to a smaller number of linear combinations of the original set of descriptors that are called principal components (PCs). The optimum number of principal components will yield the smallest standard error of

prediction while capturing as much of the variance shown by the structural data as possible. (See generally, Kramer, R., *Chemometric Techniques for Quantitative Analysis*, Marcel Dekker, Inc., 1998)

Comprehensive Descriptors for Structural and Statistical Analysis

- 5 (CODESSA) – a set of various structural descriptors typically utilized in three dimensional quantitative structure-activity studies that includes constitutional descriptors, topological descriptors, geometrical descriptors, electrostatic descriptors, and quantum mechanical descriptors, all of which require knowledge of structure beforehand. (See , Tong et al., *J. Med. Chem.*, **39**: 380-387, 1995
10 and Collantes et al., *J. Anal. Chem.*, **68**: 2038-2043, 1996, both of which are incorporated herein by reference)

- Auto-scaling** – a method whereby the quantitative spectral information contained within each particular bin is compared for all compounds in the training set to yield an average value and a standard deviation. Then, for each bin
15 comprising the structure descriptors of a given compound, the quantitative spectral information therein is expressed as a number of standard deviations above or below the average for each bin. Autoscaling equalizes the importance of inherently weak spectral signals falling within certain bins with the importance of inherently strong spectral signals falling within certain other bins in describing a
20 set of spectrally derived structure descriptors. It may also equalize the importance of different types of spectral data in a composite of spectral data.

- Fisher-weighting** – a method whereby the quantitative spectral information in bins that are important for classifying the training set compounds into different endpoint groups, such as strong and medium binders to the estrogen receptor, are
25 enhanced. For each bin, the variance of the quantitative spectral information between the endpoint groups is divided by the variance of the quantitative spectral information within the endpoint groups. The resulting dividend becomes a weighting factor that has a magnitude larger than one when a particular bin has an important role in distinguishing the endpoint groups. Each bin is multiplied by its
30 weighting factor to yield structure descriptors that are more sensitive to subtle but significant spectral variations.

Leave-one-out (LOO) Cross-Validation - a method whereby each compound in the training set is systematically excluded from the data set, after which its endpoint value is predicted by the spectral data-activity relationship derived from the remaining compounds (See, Cramer et al., *Quant. Struct-Act. Relat.* 7: 18-25, 1998, incorporated herein by reference). Cross-validation is useful for judging the reliability of a spectral data-activity relationship, especially where a validation set of compounds is not available.

EXAMPLES

The following examples further illustrate the QSAR methods of the present invention. The methods utilize spectral data as structure descriptors and correlate the spectral data with specific biological, chemical, or physical endpoints without the need to assign spectral features to their corresponding structural elements. In recognition of the advantages presented by using spectral data as structure descriptors, a correlation provided by the methods described herein is termed a Spectral Data Activity Relationship (SDAR).

In various embodiments, experimental and/or calculated molecular spectral data may be used as structure descriptors. If experimental spectral data is used in a particular embodiment, there is no need to actually know the molecular structures beforehand. Calculated molecular spectral data may be utilized as a surrogate for experimental data in some embodiments. In other embodiments, calculated molecular spectral data may be useful, in combination with an SDAR derived from experimental spectral data, for screening compounds that are created as part of a computer generated combinatorial library. In yet other embodiments, calculated molecular spectral data may be generated for compounds for which no experimental spectral data is available and used to establish an SDAR or to screen those compounds for a particular property.

In some embodiments, spectral data is used as a set of descriptors, for example descriptors of molecular structure. The pattern of the spectrum is determined, for example by segmenting the spectral data into portions covering particular spectral regions (e.g., ranges of frequency, wavelength, chemical shift,

mass to charge ratio, and the like). The number and/or the intensity of the spectral signals within each segmented region may serve as the structure descriptors. The spectral data may be of one type or be a composite of several types, including NMR, MS, IR, UV-Vis, Fluorescence, and Phosphorescence. Spectral data of a particular type may be utilized in its entirety or in part. While spectral data are often used to elucidate the structure of the compound that yields them, the information contained in the spectra may be used in some embodiments without the need to interpret the spectra. Furthermore, the spectral data may be used in certain embodiments without the need to know the structures of the compounds beforehand. Segmented spectral data is particularly amenable to encryption for secure analysis.

In some embodiments, ^{13}C NMR, EI MS, and IR spectra may be combined to yield a set of structure descriptors. Examples of these types of spectra for the compound bisphenol A are shown in Figs. 1(a), 1(b), and 1(c) respectively. Each type of spectrum contributes information regarding the structure and electronic properties of a molecule. For example, in the ^{13}C NMR spectrum shown in Fig. 1(a), the peaks reflect the number of different types of carbon atoms and the electronic environment of each type of carbon atom in bisphenol A. Peaks exhibiting the largest δ (chemical shift) values correspond to carbon atoms of a type that are the least shielded from the applied magnetic field by the electrons of the molecule. The EI MS spectrum shown in Fig. 1(b) reveals the molecular mass ($m/z = 228$, 228 amu) of bisphenol A and the mass of the fragments into which the molecule breaks up when the electrons impacting the molecule deliver enough energy to break bonds. For example, the most prominent peak in the mass spectrum shown in Fig. 1 (b) appears at a m/z of 213. The peak at $m/z=213$ corresponds to the fragment formed by the loss of a methyl group (CH_3 , $m/z = 15$) from bisphenol A (i.e. $228-213 = 15$). The mass spectral peaks thus reflect the fragments of structure that comprise the molecule.

The IR spectrum shown in Fig. 1(c) reveals the various types of bonds that are present in bisphenol A. For example the peak that appears at 3368 cm^{-1} in the IR spectrum corresponds to the two oxygen-hydrogen bonds found in bisphenol A.

While the ^{13}C NMR, EI MS and IR spectral features have been discussed with reference to the structure of bisphenol A to illustrate the types of structural information contained in these spectra, it should be noted that in some embodiments the structure of the molecule responsible for the features need not be known and no attempt need be made to assign particular spectral features to corresponding structural features.

The spectral data from Figs. 1(a), 1(b) and 1(c) are compiled in tabular form in Figs. 2(a), 2(b) and 2(c), respectively, as tables of spectral frequencies and relative intensities. Such compilations are typical of the output of modern digital spectrometers. The data as presented in these tables may be used to construct a set of molecular structure descriptors for bisphenol A and similar sets of data may be used to generate structure descriptors for any number of compounds.

To construct the set of structure descriptors shown in Fig. 3, each type of spectrum is segmented into spectral "bins" and the information content of each bin becomes a separate structure descriptor. Fig. 3 shows a set of raw structure descriptors for bisphenol A derived, for illustrative purposes only, from the spectral data as presented in the tables of Figs 2(a), 2(b) and 2(c). It is possible to include more information from the spectra than considered here, such as the intensity of IR absorption at frequencies that do not correspond to a distinct peak. Each bin in this case corresponds to a particular integer m/z ratio for the EI MS data, a range of chemical shift frequencies for the ^{13}C NMR data, and a range of wavenumber frequencies for the IR data. Moreover, each bin contains the relative intensity of the spectral feature(s) falling within a range covered by a particular bin. In this example, bins 1 to 49 were not used (because such low mass ions are common and may also result from air contamination), bins 50 to 549 correspond to m/z ratios from 50 to 549, bins 550 to 770 correspond to 1 ppm segments of the ^{13}C NMR spectrum over the range of 0 to 221 ppm, and bins 771 to 1120 correspond to 10 cm^{-1} segments of the IR spectrum over the range from 4000 cm^{-1} to 500 cm^{-1} . Those spectral bins that do not contain spectral data are given a value of zero and are not shown in Fig. 3. Other regions of the spectra can be selected to correspond to the bins in other embodiments of the invention, which is not

limited to the specific examples recited herein. In addition, the spectra can be divided into different segments, instead of the integer segments that are merely given as examples.

Tracking specific spectral data from the various spectra to the set of structure descriptors in Fig. 3 helps to illustrate how, in some embodiments, the spectral data is utilized to construct the structure descriptors. For example, the intensity of the peak at $m/z=213$ in the mass spectrum appears in bin number 213. The intensity of the peak appearing at 154.2 ppm in the ^{13}C NMR spectrum is in bin number 704 because the peak falls within the range of 154 ppm to 155 ppm which corresponds to bin number 704. The peak appearing at 3368 cm^{-1} in the IR spectrum falls within the frequency range 3370 cm^{-1} to 3360 cm^{-1} and thus the intensity of the peak appears in bin number 834 since this bin corresponds to this particular frequency range. These three entries are highlighted and shown in bold in Fig. 3.

The segmented spectral data structure descriptor set for bisphenol A shown in Fig. 3 may be altered in some embodiments to normalize the data from each type of spectrum to yield structure descriptors of similar magnitude from each type of spectral data. For example, in order to normalize the IR data and ^{13}C NMR data to the greatest value in the MS data (100), they might be expressed as the number of peaks falling within a particular bin's range, multiplied by 100. The 21 appearing in bin 834 would thus be replaced with the value 100 as would the 411 appearing in bin 704, because in both cases only one peak appears in the spectral ranges corresponding to these bins.

In other embodiments, UV-Vis, fluorescence, and phosphorescence spectral data is segmented into bins alone or in combination with each other and with NMR, MS, and IR data.

In addition to not requiring structures beforehand, experimental spectral data as used in some embodiments offers several advantages over calculated spectral data. Most experimental spectral data reflects the quantum mechanical properties of molecules as determined by both structure and solvation. However, mass spectra do not reflect solvation. Experimental spectral data also avoids the

difficulty and approximations inherent in deriving structure descriptors from quantum mechanical or electrostatic potential calculations, which are especially inaccurate when the calculations attempt to include solvation effects. Experimental spectral data may be reflective of the solution conformations that are responsible for a particular endpoint property as well as the role molecular flexibility plays in determining the endpoint property. Moreover, experimental spectral data reflect specific structural features and are therefore preferable to bulk structure descriptors such as partition coefficients that lack structural specificity. Additionally, experimental spectral data is already available for many compounds in various databases.

In some embodiments, segmented spectral data is further pre-treated before being combined with endpoint data and subjected to statistical pattern recognition or artificial intelligence based pattern recognition to extract an SDAR. Fig. 4 summarizes, in flowchart form, a particular embodiment of the steps that may be taken to establish an SDAR from spectral data. These steps and others may be performed using a computer system.

In Fig. 4, spectral data is obtained at 20, and then the spectral data is segmented into bins at 22, which in some embodiments is carried out in a manner similar to that described above with reference to Figures 1, 2, and 3. The process of segregation is repeated for the spectral data of each of the compounds in the training set. The spectral data for each of the compounds of the training set can be segmented into a matrix of bins that represent the training set. These segmented spectral data sets 24 comprise the structure descriptors for the training set compounds. Patterns in the matrix of bins can then be determined, for example by pattern recognition software.

In some embodiments, the segmented spectral data 24 is pretreated by auto-scaling 26 prior to pattern recognition, to produce auto-scaled, segmented spectral data 28. Auto-scaling is a method whereby the quantitative spectral information contained within each particular bin is averaged over all the compounds in the training set to yield an average value and a standard deviation for each bin. Then, for each bin comprising the structure descriptors of a given compound, the

quantitative spectral information therein is expressed as the number of standard deviations above or below the average value for that bin. This reduces the variation in numerical magnitude between the bins in the segmented spectral data. Auto-scaling thus helps equalize the importance of inherently weak spectral signals that fall within the spectral range of certain bins with the importance of inherently strong spectral signals that fall within the spectral range of other bins.

In some embodiments, the segmented spectral data is further pre-treated by Fisher-weighting 30 the auto-scaled data 28 to improve the ability of pattern recognition algorithms to discern which bins (and the spectral information corresponding to the spectral range covered by the bin) are most important for classifying compounds into two or more endpoint classes (such as strong and weak estrogen receptor binders). An endpoint class is assigned to each compound based upon its endpoint value. For example, compounds could be classified into two endpoint classes, those compounds that have an endpoint value above a certain number and those that have an endpoint value equal to or below a certain number. For each bin of the segmented spectral data 24 the variance in value between the endpoint classes for the bin is divided by the variance in value within the endpoint classes for the bin to yield a weighting factor. The weighting factor has a magnitude larger than one when a particular bin has an important role in distinguishing the endpoint classes.

Fisher-weighting 30 quantifies the tendency for certain bins (and the spectral range corresponding to the bin) to be more helpful than others in deciding the endpoint class of a compound. For example, consider a particular bin in which the value in the bin is always large for the compounds belonging to one endpoint class (such as strong estrogen receptor binders), and always small for the compounds of a second endpoint class (such as weak estrogen receptor binders). Knowing that the spectral data for a compound yields a large value in that particular bin helps to classify the compound into the first endpoint class. If on the other hand, the value found in the bin is always large for compounds of both classes, it reveals nothing about the endpoint class of the compound to know that it has a large value in the bin. In the first case, the variance between the classes

would be larger than the variance within each class and the weighting factor will be greater than one. Conversely, in the second case, the variance between the classes will be about equal to the variance within the class, making the weighting factor equal to about one. Each bin of the segmented spectral data is then
5 multiplied by its weighting factor to yield a set of structure descriptors 32 that emphasize the bins most important for deciding the class of a compound. Such data is more easily treated by pattern-recognition analysis.

Pattern recognition 34 is used to establish an SDAR 36, by correlating the segmented (and optionally pretreated) spectral data with the endpoint data for the
10 compounds in the training set. In particular embodiments, the compounds are classified into two or more endpoint classes (e.g. strong versus weak estrogen receptor binders) according to their relative endpoint values. The pattern recognition then determines any segmented spectral features (i.e., the bins) that are characteristic of the compounds falling into each of the classes. If Fisher-
15 weighting is performed prior to pattern recognition, the endpoint classification scheme used for Fisher-weighting may be retained during pattern recognition. A simplified example is that a bin with a strong signal in all the compounds of the test set that have strong estrogen receptor binding would be a bin that would be considered as predictive of strong estrogen receptor binding if a test compound
20 also exhibited a strong signal in that bin. Multiple such signals would, however, usually be taken into account by the pattern recognition software when determining spectral patterns that are associated with the endpoint.

Pattern recognition to establish an SDAR 36 may be accomplished by statistical methods or artificial intelligence methods. Underlying these methods is
25 the idea that if a particular bin corresponds to a spectral signal exhibited by only the compounds of a particular class (e.g. strong estrogen receptor binders), the bin will bias strongly toward that class. The extent to which a particular bin biases toward a particular class depends upon whether or not compounds from all classes (e.g. strong and weak estrogen receptor binders) exhibit spectral signals
30 corresponding to the bin. For example, if half of the compounds in one class show a signal that falls within a particular bin and slightly more than half of the

compounds in another class show a signal in the same bin, the bin may only bias slightly toward the latter class. Statistical and artificial intelligence methods attempt to quantify these biases and provide a basis for classifying compounds to an endpoint class according to their segmented spectral data.

5 Once quantified, the relationship between the segmented spectral data and the endpoint may be visualized in various ways, depending upon the particular software package utilized for pattern recognition. One way to visualize the extent to which individual bins of segmented spectral data bias toward endpoint classes is with a canonical variate factor plot. An example of a canonical variate factor plot
10 is shown in Fig. 6, which is discussed further with respect to Example 1. The SDAR model used to generate Fig. 6 was based upon two endpoint classes. The peaks pointing in one direction bias toward one class (strong estrogen receptor binding) and the peaks pointing in the other direction bias toward the other class (moderate estrogen receptor binding). The length of the peaks in Fig. 6
15 correspond to how strongly the peaks bias toward a particular endpoint class. For example, the peak appearing in bin 585 of Fig. 6 strongly biases toward the class of strong binders while the peak appearing in bin 579 of Fig. 6 biases strongly toward the other class of moderate binders.

 An SDAR, once established by pattern recognition, may be used in some
20 embodiments to quantitatively predict the endpoint value and endpoint class of any compound according to its segmented spectral data. In a particular embodiment, segmented spectral data of the type utilized to establish the SDAR is considered along with the canonical variate function to yield a prediction of endpoint class. For example, in reference to Fig. 6, a compound may be qualitatively predicted to
25 belong to the class indicated by the upward pointing peaks if it exhibits a large number of spectral features falling into the upward biasing bins. The SDAR quantifies the prediction by considering not only the number of spectral features that fall into upward or downward biasing bins, but the strength with which each bin biases toward a particular endpoint class.

30 One example of a way to visualize the ability of an SDAR to correctly classify compounds into their endpoint class is with a discriminant function plot.

An example of a discriminant function plot is given in Figure 5, which is discussed in more detail with respect to Example 1. This discriminant function plot is also based upon the same two class SDAR used to generate the canonical variate function of Fig. 6. The grey and white boxes correspond to the two endpoint classes (e.g. strong and moderate estrogen receptor binding) into which the training set data was divided. The aggregation of grey boxes and the aggregation of the white boxes, as well as the separation of the two aggregates, illustrate the ability of the SDAR to discriminate between the two endpoint classes. If the SDAR had not found a correlation between the segmented spectral data and the endpoint classes, the discriminant function would have exhibited a greater mixing of the grey and white boxes.

Unlike methods using spectral data that is referenced to a parent compound's spectral data, the methods of the present invention can in certain embodiments reveal the importance of particular structural features in determining the endpoint value without requiring knowledge of the identity of those structural features beforehand. For example, if a particular bin is identified by the canonical variates as showing a bias toward strong anti-tumor activity, subsequent elucidation of the structure responsible for the signal occupying that particular bin is possible. Furthermore, spectral data as used in some embodiments is well suited for examining structure activity relationships for compounds of diverse structures where several different groupings of atoms may elicit similar quantum mechanical features that are important to a particular endpoint property.

Another benefit of using spectral data to establish an SDAR according to the methods of the present invention is the flexibility it offers. Once spectral data are gathered for a set of compounds, an SDAR may be generated with reference to a multitude of biological, chemical, or physical endpoints for which data is
5 available. For instance, the same set of spectrally derived structure descriptors for a set of compounds may be utilized with toxicity data and antibacterial activity data for these compounds to establish two separate SDARs for the compounds, one for toxicity and the other for antibacterial activity.

Because the experimentally based SDAR method does not require input of a
10 compound's structure, the method yields accurate results without computer modeling of a compound's steric or electrostatic properties. The method also removes the need to break the compound into secondary structural motifs or assume a particular alignment for how a set of molecules will bind to another molecule.

15 Methods according to the invention additionally provide rapid and inexpensive approaches to screen compounds for selected activities or properties. Screening using these methods does not require that time and effort be expended in trying to elucidate the structure of compounds beforehand, like screening using 3D-QSAR. Screening according to the methods of the present invention, by
20 measuring a compound's spectral data and inputting it into an SDAR, is much more rapid than screening with bioassays. Furthermore, once obtained, the same spectral data for a compound may be utilized in multiple SDARs to predict various properties of the molecule. Additionally, the methods may also be used to screen mixtures or fractions of compounds for spectral features associated with a
25 particular property, thereby obviating the need to spend time and money isolating compounds that show no promise of having the particular property. The methods may therefore be extremely useful, for example, in combination with experimentally generated combinatorial libraries of compounds.

When implemented in a real-time web-based computer environment, the
30 methods of the present invention may provide investigators with a means for rapidly and securely estimating the molecular activities of proprietary molecular

structures. Spectral data as utilized in the present invention may be generated for any number of compounds and submitted by an investigator, either directly or as a segmented set of descriptors, to a central location containing the SDAR software. The spectral data or segmented spectral data may be encrypted by the user and
5 securely transmitted to the central location. Because of the speed at which pattern recognition software can make predictions based upon an SDAR, a submission of spectral data and retrieval of a prediction based on an SDAR can take place in real time over the worldwide web without the spectral data being stored at the central location. Alternatively, the SDAR and the pattern recognition software may be
10 implemented on a single computer in an investigator's laboratory. In either case, there is no need to reveal the structure of a proprietary compound to obtain a prediction. The ability to provide spectral data in a pre-segmented and encrypted form that is difficult to use for structure elucidation purposes (i.e., industrial espionage) may also encourage investigators to submit their spectral and endpoint
15 data permanently to a central location for use as part of larger training sets that may lead to more reliable SDARs.

It will be understood, however, that many variations and modifications of the methods can be made while remaining within the scope and spirit of the present invention. For example, segmentation of spectral data into bins need not occur,
20 but is only one example of how spectral patterns may be observed and compared. Other approaches for comparing spectral patterns could also be used, for example artificial neural networks that may not require gross segmentation of the spectral data per se (image analysis).

25 **Example 1- Use of ^{13}C NMR and EI Mass Spectrometric Data to Produce a Predictive Model of Estrogen Receptor Binding Activity.**

The Food Quality Protection Act, passed in 1996, mandates that the Environmental Protection Agency (EPA) develop screening and testing procedures for endocrine disrupting chemicals (EDCs). An EDC is defined as "an exogenous
30 agent that interferes with the production, release, transport, metabolism, binding, action, or elimination of natural hormones in the body responsible for the

maintenance of homeostasis and the regulation of developmental processes.”

Estrogenic compounds represent a significant subset of the EDCs to be tested.

Many of these compounds can be screened by determining how strongly the compounds bind to estrogen receptors. Thus there is a need to develop

5 inexpensive and rapid methods to screen thousands of potential compounds.

SDAR according to the invention meets this need.

The ^{13}C nuclear magnetic resonance (NMR) spectrum of a compound contains frequencies that correspond directly to the quantum mechanical properties of the molecule and depend largely on the electrostatic features and geometry, including the stereochemical configuration, of the molecule. Furthermore, a
10 solution spectrum, like NMR, inherently reflects the effects of solvation on the quantum mechanical properties. Electron-impact mass spectral (EI MS) data provide a mass-size description of molecular substructures (and possibly the whole molecule) as well as information about the strength of bonds between the atoms of
15 the molecules. The ^{13}C NMR and EI mass spectral data thus represent sets of quantum mechanical structure descriptors that reflect the electrostatic and steric properties of a molecule. Such experimental data are readily attainable and often already available.

Two Spectral Data-Activity Relationship (SDAR) models were established,
20 based on ^{13}C NMR and EI MS spectra, for 30 estrogenic chemicals whose relative binding affinities (RBA) for the alpha ($\text{ER}\alpha$) and beta ($\text{ER}\beta$) estrogen receptors are available. The SDAR models classified the 30 compounds into two endpoint classes, those with strong binding affinities for the estrogen receptors and those with medium binding affinities for the estrogen receptors. The SDAR based on
25 ^{13}C NMR data alone yielded a leave-one-out cross-validation of 90% and the SDAR based on a composite of ^{13}C NMR and EI MS data yielded a cross-validation of 83%. Two compounds that were classified incorrectly in both SDAR models were in a transition zone between these classifications. Both SDAR models correctly classified 25 compounds. The models provided rapid, reliable,
30 and simple ways to predict whether a compound binds to the estrogen receptor.

Procedure

The estrogenic relative binding affinities (RBAs) of the 30 compounds were obtained from previous publications (Kuiper et al., *Endocrinology* **138**:863-870, 1997; and Tong et al., *Endocrinology*, **138**:4022-4025, 1997). Most of the ^{13}C NMR spectrometric and EI mass spectrometric data were obtained from the Integrated Spectral Data Base System for Organic Compounds web site www.aist.go.jp/RIODB/SDBS, the *Aldrich Library of ^{13}C and ^1H FT NMR Spectra* (Pouchert and Behnke, Eds., Aldrich Chemical Company, Volumes 1-3, 1993), *Spectral Data of Steroids* (Frenkel and Marsh, eds., Thermodynamics Research Center: College Station, 1994), and the NIST MS database software version 1.6. Experimental ^{13}C NMR and EI MS data for five compounds were obtained using standard methods.

Specifically, the ^{13}C nuclear magnetic resonance (NMR) spectral analyses of 4-hydroxy-estradiol, ICI 164,384, moxestrol, and norethynodrel were performed at 75.46 MHz on a Varian Gemini 300 MHz NMR (Varian Associates, Inc., Palo Alto, CA) spectrometer operating at 301 K. The subject compounds were dissolved in CDCl_3 or DMSO (dimethyl sulfoxide). The chemical shifts were defined by denoting the CDCl_3 peak at 77.0 ppm and the DMSO peak at 39.5 ppm. The spectral width was 21,008 Hz with a 2.6-second delay time between acquisitions. The acquisition time was 0.495 seconds and the number of points acquired was 20,800.

The samples were also analyzed by direct exposure probe (DEP) mass spectrometry (MS). The mass spectrometers were operated in the electron-impact (EI) mode, with 70 V electron energy. The ion source temperature was set at 150°C. Samples, in solution, were applied to the rhenium wire of the DEP and the solvent was allowed to evaporate before the analysis was begun. MS data were collected until the current of the DEP exceeded 500 mA. Norethynodrel, ICI 164,384, 4-hydroxy-tomoxifen, and 4-hydroxy-estradiol were analyzed on a Finnigan TSQ 700 mass spectrometer and moxestrol was analyzed on a Finnigan 4500 mass spectrometer (Finnigan Corp., San Jose CA).

The ^{13}C NMR and EI-mass spectral data points were used as comprehensive descriptors for structural and statistical analyses (CODESSA) as in Tong et al., *J. Med. Chem.*, **39**: 380-387, 1995 and Collantes et al., *J. Anal. Chem.*, **68**: 2038-2043, 1996, both of which are incorporated herein by reference). Mass

5 spectrometric data for mass-to-charge (m/z) ratios from 50 to 549 were used directly as ordered pairs of m/z and relative intensity (maximum of 100).

Unassigned 1D ^{13}C NMR chemical shifts were segmented into bins over a 0 to 221 ppm range. The ^{13}C NMR frequencies were shifted to bins 550 to 770, so bin 550 was the ^{13}C NMR spectrum for frequencies inside 0 to 1 ppm and bin 770 was the

10 ^{13}C NMR spectrum for frequencies inside 220 to 221 ppm.

To save space, the ^{13}C NMR spectra were saved as sets of ordered pairs each consisting of the respective chemical shift frequency in ppm and the respective area under the peak. The area under the peak of a specific chemical shift frequency was first normalized to an integer. A non-degenerate frequency

15 was assigned an area of 1; a doubly degenerate frequency had an area of 2; and so forth. This was done to provide all the spectra with a similar signal-to-noise ratio and to eliminate line-width variations due to differences in NMR-instrument field strengths, shimming, temperature, pH, and solvents. The bin defined the number of significant and distinct chemical-shift peaks inside a ppm range. The optimal

20 range in this example was found to be 1 ppm to each bin. To normalize the peak intensity of the ^{13}C NMR spectroscopic data relative to the mass spectrometric data, the area of one ^{13}C NMR peak was set to be 25. The number 25 was selected with the objective of scaling the maximum value for the ^{13}C NMR data inside a bin to near 100, which is the maximum value for EI MS data. The number of bins

25 used to input the ^{13}C NMR spectra was studied by varying the width of the bins from 0.5 to 5.0 ppm. Again, the optimal ppm range for this study was found to be 1.0 ppm. Values were normalized to a maximum of 100 prior to pattern-recognition analysis.

The relative binding affinity (RBA) to the estrogen receptor was defined as

30 the ratio of the molar concentration of 17- β -estradiol to the competing compound required to decrease the receptor-bound 17- β -estradiol by 50%, multiplied by 100.

Thus, 17- β -estradiol has an RBA of 100 and a log (RBA) of 2.0. Strong binders to the estrogen receptor were classified as having a log (RBA) over 0.0, and medium binders to the estrogen receptor were classified as having a log (RBA) of less than or equal to 0.0. There were 17 strong binders and 13 medium binders in the training set. No weak estrogen-receptor-binding compounds (i.e., those with a log (RBA) of less than or equal to -3) were included. SDAR models that include more compounds, including weak-binding compounds, in the training set are better able to accurately predict strong estrogen receptor binding (see example 2).

The analysis of the SDAR model was performed by the leave-one-out (LOO) cross-validation procedure in which each compound was systematically excluded from the training set and its relative binding activity class predicted by the model (Cramer et al., *Quant. Struct.-Act. Relat.*, 7:18-25, 1988, incorporated by reference).

The pattern-recognition software used was RESolve Version 1.2 (Colorado School of Mines, Boulder, CO). The ^{13}C NMR and EI MS spectroscopic data for all 30 compounds were input as text files into a computer programmed with the software. The spectroscopic data was then auto-scaled and Fisher-weighted prior to principal component analysis (PCA). The discriminant analysis was based on canonical variate vectors. Leave-one-out (LOO) cross-validation was used to maximize the size of the training set.

Autoscaling compared the quantitative response at each mass spectral m/z bin or NMR chemical shift bin to all the other bins in the comparison set. An average value (with standard deviation) was calculated for each bin. Then, for each spectrum, the quantitative response at each bin was expressed as the number of standard deviations above or below the respective average. This data-pretreatment step equalized the weight of consistent variance of signals with inherently small magnitudes (25 units for the NMR bin 558 representing a single methyl carbon) to those signals with large magnitudes (130,000 area counts at m/z bin 91, possibly arising from a tropylium ion fragment). Autoscaling automatically compensates for gross magnitude variations. It was particularly helpful in this example in which two completely different types of analytical

spectra were formed into a composite set of structure descriptors that are representative of molecular characteristics.

Fisher weighting was used to emphasize those spectral characteristics important in distinguishing defined endpoint groups. For each mass spectral m/z or NMR bin, the variance between groups was divided by the variance within groups. The resulting dividend became a weighting factor with a magnitude larger than one when the particular m/z or bin has an important role in distinguishing groups. Fisher weighting all of the raw spectra before pattern recognition increased the power of discriminant analysis to classify spectra correctly. In this example, Fisher weighting de-emphasized the relative importance of irrelevant spectral information (such as the NMR signals from carbon atoms not important to binding to the estrogen receptor).

Results

Based only on ^{13}C NMR spectroscopic data, the statistical pattern-recognition program with 8 principal components (PCs) had 81.1% of the total variance and a cross validation of 90%. The mathematical mode of significance was 91.9% at 1, with 28 degrees of freedom. The Wilks discriminant criterion was 99.99%. Table 1 lists the compound names, an identifying number for each compound that is used in Figs. 5 and 7, the log (RBA) alpha and log (RBA) beta values, the SDAR training input endpoint class for each compound ("class input")(either S for strong binding or M for medium binding), the SDAR prediction ("class predicted NMR") from the ^{13}C NMR data alone and the SDAR prediction ("class predicted NMR and MS") from the ^{13}C NMR and EI MS data. Fig. 5 shows the discriminant function for SDAR derived from the ^{13}C NMR data alone which illustrates the ability of the SDAR to discriminate between the two endpoint classes in the training set. Compounds denoted with a white background exhibit strong RBAs, and compounds with a gray background exhibit medium RBAs. Fig. 5 shows that the SDAR yielded a large separation between the 15 strong (white rectangles on right side) and the 9 medium binders (gray rectangles on the left side); 4 strong and 2 medium RBAs are in a transition zone (middle)

between strong and medium RBAs. Table 1 also shows that 27 of the 30 compounds are correctly group-predicted using only, ^{13}C NMR data. 3β -androstanediol, 2-hydroxy-estradiol and hexestrol are incorrectly predicted to have a medium RBA using 8 principal components. Only 3β -androstanediol and
5 hexestrol are incorrectly predicted to have a medium RBA using the canonical variate function. Strong binding 3β -androstanediol is most likely incorrectly predicted because the compound is similar to the medium RBA 3α -androstanediol. Repeating the analysis using a larger training set (Example 2) eliminated even this confusion.

TABLE 1

| COMPOUND | Number | Log(RBA) alpha | Log(RBA) beta | Class Input | Class Predicted NMR | Class Predicted NMR + MS |
|-----------------------------|--------|-------------------|------------------|----------------|---------------------------|--------------------------------|
| Coumestrol | 1 | 1.97 | 2.27 | S | S | S |
| 3 α -Androstenediol | 2 | -1.15 | -0.52 | M | M | S |
| 3 β -Androstenediol | 3 | 0.48 | 0.85 | S | M | M |
| 4-OH-Estradiol | 4 | 1.11 | 0.85 | S | S | S |
| 17 α -Estradiol | 5 | 1.76 | 1.04 | S | S | S |
| 17 β -Estradiol | 6 | 2.00 | 2.00 | S | S | S |
| Bisphenol A | 7 | -1.30 | -0.48 | M | M | M |
| β -Zearanol | 8 | -1.20 | -1.15 | M | M | M |
| Clomifene | 9 | 1.40 | 1.08 | S | S | S |
| 2-OH-Estradiol | 10 | 0.85 | 1.04 | S | M | S |
| Dehydroepiandrosterone | 11 | -1.40 | -1.15 | M | M | M |
| Diethystibesterol | 12 | 2.67 | 2.47 | S | S | S |
| Dienestrol | 13 | 2.35 | 2.61 | S | S | S |
| Dihydrotestosterone | 14 | -1.30 | -0.77 | M | M | M |
| Estriol | 15 | 1.15 | 1.32 | S | S | S |
| Estrone | 16 | 1.78 | 1.57 | S | S | S |
| Genistein | 17 | 0.70 | 1.56 | S | S | S |
| Hexestrol | 18 | 2.48 | 2.37 | S | M | M |
| ICI 164,384 | 19 | 1.93 | 2.22 | S | S | S |
| Methoxychlor | 20 | -2.00 | -0.89 | M | M | M |
| Moxestrol | 21 | 1.63 | 0.70 | S | S | S |
| Nafoxidine | 22 | 1.64 | 1.20 | S | S | S |
| Nandrolone | 23 | -2.00 | -0.64 | M | M | M |
| Norethindrone | 24 | -1.15 | -2.00 | M | M | M |
| Norethynodrel | 25 | -0.16 | -0.66 | M | M | M |
| Progesterone | 26 | -3.50 | -3.50 | M | M | M |
| Tamoxifen | 27 | 0.85 | 0.78 | S | S | S |
| Testosterone | 28 | -2.10 | -2.10 | M | M | M |
| 5 α -Androstenedione | 29 | <-2.0 | <-2.0 | M | M | S |
| 5 β -Androstenedione | 30 | <-2.0 | <-2.0 | M | M | S |

- 5 Fig. 6 shows the canonical variate function for the pattern recognition obtained with the ^{13}C NMR data only. The positive (upward-pointing) peaks in Fig. 6 correspond to bins that bias toward a strong RBA for binding to the estrogen receptor and negative (downwardly-pointing) peaks correspond to bins that bias toward a medium RBA. In Fig. 6, m/z bins 550 to 770 refer to the ^{13}C
- 10 NMR data from 0 ppm to 1 ppm for bin 550 to data from 221 ppm to 222 ppm for bin 770. The canonical variate function reveals that the aliphatic CH_2 bins 580 to 585 (30 to 35 ppm) have a bias toward medium RBA. It also reveals that the

methyl CH₃ bins 558 and 566 (8 and 16 ppm, respectively) have a bias toward strong RBA. Many of the benzyl bins 665 to 700 (115 to 150 ppm) have a bias towards strong RBA. Thus these methyl groups and benzyl atoms are indicated to be important for strong RBA.

5 Based on the composite ¹³C NMR and EI MS data, the statistical pattern-recognition program with 14 principal components (PCs) included 89.8% of the total variance and had a cross validation of 83.3%. The mathematical mode of significance was 97.6% at 1, with 28 degrees of freedom. The Wilks discriminant criterion was 99.89%. Fig. 7 shows the discriminant function for the composite
10 ¹³C NMR and EI MS data. Compounds with a white background are strong RBAs and compounds with a gray background are medium RBAs. Fig. 7 shows a large separation between 15 strong (left) and 9 medium (right) binders. There are 2 strong and 4 medium binders in the transition zone between the strong and medium binders. Table 1 shows that 25 of the 30 compounds are correctly predicted using
15 the composite ¹³C NMR and EI/MS data. Only 3β-androstanediol and hexestrol are incorrectly predicted using the canonical variate function. Again, the strong binding 3β-androstanediol is most likely predicted incorrectly because it is similar to the medium binding 3α-androstanediol.

Fig. 8(a) shows the canonical variate function used in the pattern
20 recognition of the EI MS data. Fig. 8(b) shows the canonical variate function used in the pattern recognition of the ¹³C NMR data. The negative peaks in Figs 8(a) and 8(b) correspond to bins that bias toward a strong relative binder and positive peaks correspond to bins that bias toward a medium relative binder, to the estrogen receptor. In Figs. 8(a) and 8(b), the label "m/z bins 50 to 550" refers to the EI
25 MS data, whereas "m/z bins 550 to 770" refers to the ¹³C NMR data. In Fig. 8(a), the mass canonical variate is split evenly into bins that bias strong and medium binding. Many of the canonical variate bins that showed bias in Fig. 2 are also present in Fig. 8(b), but they are pointing in the opposite direction. The opposite directions found in Fig. 6 and 8(b) are insignificant technically, arising from the
30 pattern-recognition program's arbitrary choice of left and right in the corresponding canonical variate score plot.

Only 3β -androstanediol and hexestrol are incorrectly predicted in both SDAR models. 3β -androstanediol is closest to 3α -androstanediol, 5α -androstanedione, and 5β -androstanedione, all three of which have medium RBAs. Note however, 3β -androstanediol has been inconsistently identified in the literature
5 as a medium estrogen-receptor binder and a strong estrogen-receptor binder (see Miksicek, *J. Steroid. Biochem. Mol. Biol.* **49**:153-160, 1994 and Kuiper et al., *Endocrinology* **138**:863-870, 1997, respectively).

This example illustrates that, using the present method, compounds can be classified by biological activity based on similarities in the spectral patterns of the
10 test compounds to the spectral patterns of the training set. In particular, additional test compounds can be classified according to their expected biological activity by obtaining the corresponding spectral pattern(s) of the test compound (e.g. NMR and MS patterns). Expected biological activity of the test compound can be predicted by detecting similarities between the spectral pattern of the test
15 compound and the spectral patterns of compounds in the training set that are associated with a known biological activity. Similarities in patterns of the canonical variates associated with an endpoint (e.g. a biological activity such as estrogen receptor binding) derived for the training compounds with patterns in the test compound's segmented spectral data, are then used to predict whether the test
20 compound will share the endpoint.

Example 2 – Expanded Training Set for Estrogen Receptor Binding SDAR

An expanded training set of 119 compounds of varying estrogen receptor
25 binding affinities was utilized to create an SDAR model from ^{13}C NMR spectral data and to create an SDAR model from a composite of ^{13}C NMR spectral data and EI MS spectral data. The training set included weak estrogen receptor binders in addition to strong and medium estrogen receptor binders. For this model, strong estrogen receptor binders were defined as having $\log(\text{RBA}) > -0.25$ and weak
30 estrogen receptor binders were defined as having $\log(\text{RBA}) < -3.0$. Medium estrogen receptor binders were defined as having $\log(\text{RBA})$ in between. The

compound names, the input class ("Class"), the predicted endpoint class based upon ^{13}C NMR only ("Class predicted NMR"), and the predicted endpoint class based upon ^{13}C NMR plus EI MS ("Class predicted NMR/MS") for the training set are listed below in Table 2. Except for the increased number of compounds and a change in the mass spectral range utilized, the analysis was performed in the same manner as the analysis in Example 1.

Table 2

| Compound | Log(RBA) | Class | Class predicted NMR | Class predicted NMR/MS |
|---------------------------------------|----------|-------|---------------------|------------------------|
| 1,4-Diphenyl-1,3-butadiene | | W | W | W |
| 1,6-Dimethylnaphthalene | | W | W | W |
| 1,8-Octanediol | | W | M | M |
| 2,2'-Dihydroxy-4-methoxybenzophenone | | W | W | W |
| 2,2'-Methylenebis (4-chlorophenol) | -2.45 | M | M | M |
| 2,4,5-T | | W | W | W |
| 2,4-D | | W | W | W |
| 2,4-Dihydroxybenzophenone | -2.61 | W | M | M |
| 2-Chloro-4-methylphenol | -3.66 | W | W | W |
| 2-Chlorophenol | -3.67 | W | W | W |
| 2-Furaldehyde | | W | W | W |
| 2-OH-Estradiol * | | S | S | S |
| 2-Phenylphenol | | W | W | W |
| 2-sec-Butylphenol | -3.54 | W | W | M |
| 3-Phenylphenol | -3.44 | W | W | W |
| 4,4'-Dihydroxybenzophenone | -2.46 | M | W | M |
| 4,4'-Methylenebis(N,N-dimethyl) | | W | M | W |
| 4,4'-Methylenedianiline | | W | W | W |
| 4,4'-Methylene(2,6-ditertbutylphenol) | | W | W | W |
| 4,4'-Sulfonylphenol | -3.07 | W | W | W |
| 4-Benzyloxyphenol | -3.44 | W | W | W |
| 4-Chloro-2-methylphenol | -3.67 | W | W | W |
| 4-Chloro-3-methylphenol | -3.38 | W | W | W |

| | | | | |
|---|-------|---|---|---|
| 4-Ethylphenol | -4.17 | W | W | W |
| 4-Nonylphenol | -1.45 | M | M | M |
| 4-OH-Estradiol * | | S | S | S |
| 4-OH-Tamoxifen | 2.24 | S | S | W |
| 4-Phenylphenol | -3.04 | W | W | W |
| 4-Stilbenol | | W | W | S |
| 4-tert-Amylphenol | -3.26 | W | W | W |
| 4-tert-Butylphenol | -3.61 | W | W | W |
| 4-tert-Octylphenol | -1.82 | M | W | W |
| 5 α -Androstane-3 α ,17 β -diol | -2.67 | M | W | M |
| 5 α -Androstane-3 β ,17 β -diol | -0.92 | M | M | M |
| Aldrin | | W | W | W |
| Apigenin | -1.61 | M | M | M |
| Aurin | -1.49 | M | M | W |
| Benzylalcohol | | W | W | W |
| Benzylbutyl phthalate | | W | W | W |
| BiochaninA | -2.37 | M | M | M |
| BIS (2-ethylhexyl) adipate | | W | M | M |
| BIS (2-ethylhexyl) phthalate | | W | W | W |
| BIS (2-hydroxyphenyl)-methane | | W | W | W |
| BIS (4-hydroxyphenyl)-methane | -3.02 | W | W | W |
| Bisphenol A | -2.11 | M | M | M |
| Bisphenol B | -1.07 | M | S | M |
| Butyl-4-aminobenzoate | | W | W | W |
| n-Butylbenzene | | W | W | W |
| Caffeine | | W | M | W |
| Catechin | | W | S | S |
| Chalcone | -2.82 | W | S | W |
| Cholesterol | | W | W | W |
| Chrysene | | W | W | W |
| Chrysin | | W | M | M |
| Cineole | | W | W | W |
| Cinnamic acid | | W | W | W |
| Clomiphene | -0.14 | S | W | W |
| Corticosterone | | W | W | S |
| Coumestrol | -0.05 | S | S | S |

| | | | | |
|---|-------|---|---|---|
| DDD-o,p' | | W | W | W |
| DDD-p,p' | | W | W | W |
| DDE-p,p' | | W | S | S |
| DDT-o,p' | -2.85 | W | W | W |
| DDT-p,p' | | W | W | W |
| DES | 2.60 | S | S | S |
| Daidzein | -1.65 | M | M | M |
| Dexamethasone | | W | S | W |
| Dibenzo-18-crown-6 | | W | W | W |
| Dibutyl phthalate | | W | W | W |
| Dieldrin | | W | S | S |
| Dienestrol | 1.57 | S | M | S |
| Diethyl phthalate | | W | W | W |
| Dihydrotestosterone | | W | M | W |
| Diisobutyl phthalate | | W | W | S |
| Dimethyl phthalate | | W | W | W |
| Diphenolic acid | -3.13 | W | M | S |
| Dopamine | | W | W | W |
| Estra-1,3,5(10)-trien-3-ol | 1.14 | S | S | W |
| Estra-1,3,5(10)-trien-3,6 α ,17 β -triol | -0.15 | S | S | S |
| Estriol | 0.99 | S | S | S |
| Estrone | 0.86 | S | S | S |
| Ethylcinnamate | | W | W | W |
| Ethynyl estradiol | 2.28 | S | S | S |
| Etiocholan-17 β -ol-3-one | | W | S | S |
| Eugenol | | W | W | W |
| Flavanone | | W | W | M |
| Flavone | | W | W | W |
| Genistein | -0.35 | M | M | M |
| Heptanal | | W | W | W |
| Hesperetin | | W | M | M |
| Hexachlorobenzene | | W | W | W |
| Hexestrol | 2.48 | S | W | W |
| Hexylalcohol | | W | W | W |
| ICI-164,384 | 1.16 | S | S | S |
| Isoeugenol | | W | W | W |

| | | | | |
|------------------------|-------|---|---|---|
| Kaempferol | -1.55 | M | M | M |
| Lindane | | W | W | S |
| Melatonin | | W | M | W |
| Mestranol | 0.35 | S | S | S |
| Methoxychlor | -3.2 | W | W | W |
| Morin | -3.35 | W | W | W |
| Moxestrol | 1.14 | S | S | S |
| Nafoxidine | -0.14 | S | S | S |
| Naringenin | -2.13 | M | W | W |
| Naringin | | W | W | W |
| Norethynodrel | -0.65 | M | M | M |
| Phenolphthalein | -1.87 | M | W | W |
| Phenol red | -3.25 | W | W | W |
| Phloretin | -1.24 | M | M | M |
| Progesterone | | W | W | W |
| Quercetin | | W | W | M |
| Suberic acid | | W | W | W |
| Tamoxifen | 0.21 | S | S | S |
| Testosterone | | W | W | W |
| Triphenylethylene | -2.78 | W | W | W |
| Triphenylphosphate | | W | W | W |
| Vanillin | | W | W | W |
| 17 α -Estradiol | 0.49 | S | S | S |
| 17 β -Estradiol | 2.0 | S | S | S |

Based only on ^{13}C NMR spectroscopic data, the statistical pattern recognition program with 22 principal components (PCs) had 88.9% of the total variance and a cross validation of 77.3 %. Figure 9 shows the discriminant function for ^{13}C NMR data of 119 compounds. Compounds that are represented by an S are strong estrogen receptor binders, compounds that are represented by an M are medium estrogen receptor binders, and compounds that are represented by a W are weak estrogen receptor binders. Figure 9 shows a clustering between the 20 strong, 19 medium, and 80 weak binders. There are cluster overlaps between the weak and strong binders and between the weak and medium cluster

region. There are a total of eight false positives and two false negatives when using LOO to predict from the 119 compound SDAR model of estrogen receptor binding using only ^{13}C NMR data.

Figure 10 shows the factors associated with the first canonical variate function for the pattern recognition of the ^{13}C NMR data. The positive peaks in Figure 10 correspond to bins that bias toward a strong RBA for binding to the estrogen receptor and negative peaks correspond to bins that bias toward a medium RBA. In Figure 10, m/z bins 550 to 770 refer to the ^{13}C NMR data from 0 ppm for bin 550 to 220 ppm for bin 770. The aliphatic CH_2 bins 580 to 585 (30 to 35 ppm) have a bias toward medium RBA. The methyl CH_3 bins 558 and 566 (8 and 16 ppm, respectively) have a bias toward strong RBA. Many of the aromatic bins 665 to 700 (115 to 150 ppm) have a bias towards strong RBA.

Based on the composite ^{13}C NMR and EI/MS data over the range $m/z = 100$ to $m/z = 549$, the statistical pattern recognition program with 22 principal components (PCs) included 80.1% of the total variance and had a cross validation of 79.0 %. Figure 11 shows the discriminant function for ^{13}C NMR data for 119 compounds. The compounds that are shown by an S exhibit strong RBAs, compounds that are shown by an M are exhibit medium RBAs, and compounds that are shown by a W exhibit weak RBAs. Figure 11 shows a similar clustering between the 20 strong, 19 medium, and 80 weak binders as seen Figure 9. There are cluster overlaps between the weak and strong binders and between the weak and medium cluster region. There are a total of nine false positives and three false negatives when using LOO to predict a compound's endpoint class from the 119 compound SDAR model of estrogen receptor binding using ^{13}C NMR and EI/MS data.

Figures 12(a) and 12(b) shows the factors associated with the first canonical variate function for the pattern recognition of the composite ^{13}C NMR and EI/MS data. The positive peaks in Figures 12(a) and 12(b) correspond to bins that bias toward a strong RBA for binding to the estrogen receptor and negative peaks correspond to bins that bias toward a medium RBA. In figure 12(a), m/z bins 100 to 549 are the EI/MS data bins and in Figure 12(b), m/z bins 550 to 770 refer to

the ^{13}C NMR data from 0 ppm for bin 550 to 220 ppm for bin 770. The aliphatic CH_2 bins 580 to 585 (30 to 35 ppm) have a bias toward medium RBA. The methyl CH_3 bins 558 and 566 (8 and 16 ppm, respectively) have a bias toward strong RBA. Many of the aromatic bins 665 to 700 (115 to 150 ppm) have a bias towards strong RBA.

In both 119 compound SDAR models of estrogen receptor binding, a possible cutoff for testing experimentally whether a molecule did bind to the estrogen receptor would be anything with a positive component 1 as depicted in figures 9 and 11. This would leave no positives untested in each model. In both models, we would test 11 false negatives and cut the number of molecules to be tested for binding to the estrogen receptor to 31 out of the original 119 compounds.

A particularly striking result of the analysis using strong, medium, and weak estrogen receptor binders in the training set is the good separation between strong estrogen receptor binders and the other compounds as seen in Figures 9 and 11. There is a marked intermixing of the medium and weak estrogen receptor binders in Figures 9 and 11 with respect to the first canonical variate function (X-axis). These results are understandable in light of the physical basis for estrogen receptor binding. Estrogen receptor binding, like most ligand-receptor binding interactions, is governed by specific intermolecular interactions between the ligand and the receptor. The specific intermolecular interactions are what distinguish strong binders from the other compounds. Conversely, there are few or no specific interactions with the receptor on which to distinguish the medium and weak estrogen receptor binders from each other.

**Example 3 - Use of ^{13}C NMR, EI MS, and Infrared Spectral Data to
Produce a Predictive Model of Biodegradability for Monocyclic Chlorobenzene
Derivatives**

5

Two recognized factors in the biodegradation of xenobiotic compounds in the environment are the physicochemical properties of the compounds themselves, and the enzymatic activities expressed by the microbial communities in the environment. The vast majority of microbial enzymatic reactions may be

10 classified into one of four categories: 1) group transfer; 2) oxidation and reduction; 3) elimination, isomerization, and rearrangement; and 4) carbon-carbon bond cleavage. The extent to which xenobiotic chemicals can serve as substrates for these enzymatic reactions is determined to a large extent by their structure and particularly by the presence of bonds that are similar to those found in the natural

15 substrates for the enzymes.

Infrared (IR) spectroscopic data may be included in an SDAR to provide a set of spectrally derived structure descriptors that include the types of bonds present in the subject molecules. Infrared data also reflect the modes and frequencies of vibration that are available to subject molecules. As chemical

20 reactions are often tied to particular vibrations of particular bonds, inclusion of the infrared data may improve the ability of an SDAR to predict reactivity.

The biodegradation data for many monocyclic chlorobenzene derivatives in sediment may be found in the Database for Environmental Fate of Chemicals (www.aist.go.jp/RIODB/dbefc). Additional data on the biodegradability is

25 published in *Biodegradation and Bioaccumulation Data of Existing Chemicals Based on the CSCL*, Japan Chemical Industry Ecology-Toxicology & Information Center (JETOC), Tokyo, Japan, 1992. The half-life period is used as the endpoint for the establishment of the SDAR and compounds are classified into two endpoint classes as readily biodegradable (R) (half-life < 30 days) and not readily

30 biodegradable (NR) (half-life > 30 days). The endpoint data for 34 chlorobenzene derivatives are given in Table 3 below.

Table 3

| Compound | Half-life (days) | Input Class |
|-----------------------------|------------------|-------------|
| Monochlorobenzene | 46.2 | NR |
| 1,2-dichlorobenzene | 36.9 | NR |
| 1,3-dichlorobenzene | 433 | NR |
| 1,4-dichlorobenzene | 385 | NR |
| 1,2,3-trichlorobenzene | 23 | R |
| 1,2,4-trichlorobenzene | 40.8 | NR |
| 1,3,5-trichlorobenzene | 35 | NR |
| 1,2,3,4,-tetrachlorobenzene | 18.2 | R |
| 1,2,3,5-tetrachlorobenzene | 18.6 | R |
| 1,2,4,5-tetrachlorobenzene | 28.8 | R |
| Pentachlorobenzene | 17.8 | R |
| Hexachlorobenzene | 27.1 | R |
| 2-chloroaniline | 175 | NR |
| 3-chloroaniline | 672 | NR |
| 4-chloroaniline | 203 | NR |
| 2,3-dichloroaniline | 54 | NR |
| 3,4-dichloroaniline | 55 | NR |
| 3,5-dichloroaniline | 149 | NR |
| 3,4,5-trichloroaniline | 145 | NR |
| 2,3,4,5-trichloroaniline | 57 | NR |
| Pentachloroaniline | 40 | NR |
| 2-chlorophenol | 6.9 | R |
| 3-chlorophenol | 29.2 | R |
| 4-chlorophenol | 11.7 | R |
| 2,3-dichlorophenol | 13.2 | R |
| 2,4-dichlorophenol | 15.3 | R |
| 2,5-dichlorophenol | 46.3 | NR |
| 2,6-dichlorophenol | 5.0 | R |

| | | |
|---------------------------|------|----|
| 3,4-dichlorophenol | 69.9 | NR |
| 3,5-dichlorophenol | 17.3 | R |
| 2,3,4-trichlorophenol | 1.8 | R |
| 3,4,5-trichlorophenol | 21.7 | R |
| 2,3,4,5-tetrachlorophenol | 6.5 | R |
| Pentachlorophenol | 2.1 | R |

Spectral data for these chlorobenzene compounds is obtained from the Integrated Spectral Data Base System for Organic Compounds web site, www.aist.go.jp/RIODB/SDBS, the *Aldrich Library of ^{13}C and ^1H FT NMR Spectra* (Poucher and Behnke, Eds., Aldrich Chemical Company, Volumes 1-3, 1993) and the NIST MS database software version 1.6. Experimental ^{13}C NMR, EI MS, and IR data is collected when spectral data is not available in a database. Experimental spectral data is collected using standard spectroscopic protocols.

The spectral data points are segmented and used in the same way as 3D-
10 QSAR uses comprehensive descriptors for structural and statistical analyses (CODESSA) (See, Tong et al., *J. Med. Chem.*, **39**: 380-387, 1995 and Collantes et al., *J. Anal. Chem.*, **68**: 2038-2043, 1996, both of which are incorporated herein by reference). Specifically, the ^{13}C NMR, EI-mass spectral, and IR data are placed into bins. Mass spectrometric data from m/z of 50 to 549 are used.
15 Unassigned 1D ^{13}C NMR chemical shifts are segmented into bins over a 0 to 221 ppm range. The ^{13}C NMR frequencies are shifted to bins 550 to 770, so bin 550 is the ^{13}C NMR spectrum for frequencies inside 0 to 1 ppm and bin 770 is the ^{13}C NMR spectrum for frequencies inside 220 to 221 ppm. The unassigned IR frequencies in the range from 4000 cm^{-1} to 600 cm^{-1} are segmented into bins of 50
20 cm^{-1} width. The IR data is shifted to bins 771 through 839 so that bin 771 holds the spectral data for the range 4000 cm^{-1} to 3950 cm^{-1} and bin 839 holds the spectral data for the range 650 cm^{-1} to 600 cm^{-1} . As an alternative, the IR data may be segmented into ranges of frequency or wavelength rather than the customary wavenumber frequencies.

To save space, the ^{13}C NMR spectra are saved as sets of ordered pairs, each consisting of the respective chemical shift frequency in ppm and the number of peaks within the frequency range. Likewise, the IR spectra are saved as ordered pairs of bin number and the number of distinct IR peaks appearing in that particular range. Frequency ranges with only one peak are first assigned an area of 1; doubly degenerate frequencies an area of 2; and so forth. This is done to provide all the spectra with similar signal-to-noise ratios and to de-emphasize line-width variations due to differences in experimental conditions. The bin thus defines the number of significant and distinct spectral features within a frequency range. Normalization of the ^{13}C NMR spectroscopic data and the IR data is accomplished by multiplying each set by a number that makes the maximum value in their respective set of bins equal to 100, which is the maximum value for EI MS data. The number of bins used to input ^{13}C NMR spectra and IR spectra may be varied to improve the SDAR. Increasing the number of bins and shrinking the frequency width of each bin provides separate bins for closely spaced spectral features that may prove important as structure descriptors for establishing a reliable SDAR.

The segmented ^{13}C NMR, EI MS, and IR spectral data along with the biodegradability data is input into the pattern recognition program RESolve Version 1.2 (Colorado School of Mines, Boulder, CO) and is auto-scaled and Fisher-weighted prior to principal component analysis (PCA). The discriminant analysis is performed based upon the canonical variate vectors. Leave-one-out (LOO) cross-validation is used to maximize the size of the training set. Spectral data for a test compound is then input and the compound judged to be readily biodegradable or not readily biodegradable based upon the presence of a majority of spectral features falling in bins that correspond to canonical variate factors from the training set that bias toward readily biodegradable or not readily biodegradable.

Alternatively, an expert system (self-learning or not) or artificial neural network may be utilized to perform the pattern recognition. Where the endpoint may be the result of multiple underlying mechanisms, such as multiple pathways of

biodegradation, expert systems or artificial neural networks may be advantageously utilized to separate compounds based upon their mode of biodegradation in addition to their biodegradability.

One example of an expert system is the MuRES system that is also part of the RESolve 1.2 software package. The MuRES package uses spectral data that is first compressed by projection onto a set of principal components. This method of compression is utilitarian, because it reduces the number of variables while maximizing the information content. MuRES uses the spectral scores that are calculated by projecting the spectral data onto a set of eigenvectors. However, MuRES may be applied directly without the compression step to data that already is overdetermined (i.e. more compounds and endpoint values than spectral data structure descriptors).

The knowledge base created by MuRES is in the form of simple binary rules. Binary rules use binary logic, logic that can be only true or false. A complex solution to a problem may be decomposed into a tree, often referred to as a classification tree, which consist of simple binary rules. The largest advantage of the expert system is that the scores do not have to be linearly separable which is an assumption required by discriminant analysis. Further details of the MuRES method are found in Harrington, *RESolve Software Manual*, Colorado School of Mines, Golden CO, which is incorporated herein by reference.

Example 4 – Use of ^{13}C NMR and UV/Visible Spectroscopic Data to Produce a Predictive Model for the Photosensitized Production of Singlet Molecular Oxygen.

Phototosensitized oxidations involving singlet oxygen, a strong oxidant, are implicated in photodynamic inactivation of viruses and cells, in phototherapy for cancer, in photocarcinogenesis and in photodegradation of dyes and polymers. Quenching of excited singlet and triplet states of many substances by ground state molecular oxygen produces singlet oxygen; the lowest electronically excited singlet state of molecular oxygen. A compilation of the quantum yields for the formation of singlet oxygen in fluid solutions for over 700 substances is available from the

Notre Dame Radiation Laboratory – Radiation Chemistry Data Center

(<http://www.rcdc.nd.edu>).

Compounds that are capable of photosensitizing the production of singlet oxygen are quite diverse and include aromatic hydrocarbons, aromatic ketones and thiones, quinones, coumarins, fluoresceins, transition metal complexes, and heterocyclics. Porphyrins and phthalocyanines are particularly important classes of compounds that are capable of producing singlet oxygen upon illumination.

To establish an SDAR useful for predicting the ability of molecules to photosensitize the production of singlet oxygen, the quantum yields for production of singlet oxygen by 20 compounds of diverse structure are obtained from the Radiation Chemistry Data Center website. The compounds are divided into two endpoint classes based upon having a high quantum yield for production of singlet oxygen (H, QY > 0.50) or a low quantum yield for production of singlet oxygen (L, QY < 0.50). Spectral data for these compounds is obtained from the resources mentioned in Examples 1 and 2 or is gleaned from other literature sources or is measured experimentally. The compounds, their singlet oxygen quantum yields, and the endpoint class for each are listed in Table 4.

Table 4

| Compound | Quantum Yield | Endpoint Class |
|--------------------|---------------|----------------|
| Acetophenone | 0.29 | L |
| Acridine | 0.83 | H |
| Anthracene | 0.83 | H |
| 9,10-Anthraquinone | 0.15 | L |
| Anthrone | 0.25 | L |
| Azulene | 0 | L |
| Benzil | 0.57 | H |
| Benzophenone | 0.29 | L |
| Biacetyl | 0.57 | H |

| | | |
|---------------------------|--------|---|
| Biphenyl | 1.0 | H |
| Flourene | 0.74 | H |
| 1-Methylindole | 0.35 | L |
| 1,3-Diphenylisobenzofuran | 0.26 | L |
| Phenalen-1-one | 0.94 | H |
| Pivalothiophenone | 1.0 | H |
| Psoralen | 0.0055 | L |
| Pyrene | 0.71 | H |
| (E)-Stilbene | 0.08 | L |
| Tetracene | 0.70 | H |
| Xanthen-9-one | 0.33 | L |

Ultraviolet-visible (UV-Vis) spectral data is used along with ^{13}C NMR spectral data in the SDAR to include a measure of the importance of molecular excited states to the production of singlet oxygen. The segmented spectral data is used as a set of descriptors in the same way that 3D-QSAR uses comprehensive descriptors for structural and statistical analyses. (CODESSA) (See , Tong et al., *J. Med. Chem.*, **39**: 380-387, 1995 and Collantes et al., *J. Anal. Chem.*, **68**: 2038-2043, 1996, both of which are incorporated herein by reference). The ^{13}C NMR spectral data is segmented into 1ppm bins over a 0 to 222ppm range. The UV-Vis data is segmented into 5 nm bins over the range of 190 nm to 900 nm. The ^{13}C NMR spectral data occupies bins 1 through 221 and the UV-Vis data occupies bins 222 through 321. To save space, the ^{13}C NMR spectral data is saved as sets of ordered pairs, each consisting of the bin number and the number of peaks with the frequency range corresponding to the bin. As described above, the ^{13}C NMR bins define the number of significant and distinct spectral features within a frequency range. The UV-Vis data is saved as sets of ordered pairs, each consisting of the bin number and the average molar absorptivity of the molecule within the wavelength range corresponding to the bin. Molar absorptivity is used rather than absorbance to correct for variations in concentration between the measured UV-Vis spectra of the compounds.

The segmented ^{13}C NMR and UV-Vis spectral data along with the quantum yields for singlet oxygen production for 20 compounds (the training set) are input as text files into the pattern recognition program RESolve Version 1.2 (Colorado School of Mines, Boulder CO). The spectroscopic data is auto-scaled and Fisher-weighted prior to principal component analysis (PCA). The discriminant analysis is performed based upon the canonical variate vectors. Leave-one-out (LOO) cross-validation is used to maximize the size of the training set and measure the ability of the SDAR to classify compounds correctly into their endpoint class. The number of bins used to input ^{13}C NMR spectral data and UV-Vis spectral data may be varied. Increasing the number of bins will shrink the frequency width of each bin and provide separate bins for closely spaced spectral features that may prove important as structure descriptors for establishing a reliable SDAR. Increasing the number of compounds in the training set may also lead to a more reliable SDAR.

Once the training set has been cross-validated and deemed reliable, test compounds are then subjected to ^{13}C -NMR and UV/Visible Spectroscopy, and these spectra are segmented into bins in the same manner as with the training set. Spectral patterns in these spectra for the test compound are then compared to the patterns for the training set, and endpoints associated with the spectral patterns of the training set are used to predict endpoints for the test compound. Where the endpoint is photosensitization of singlet oxygen, the presence or absence of spectral patterns in the training set which are associated with that endpoint are then detected to predict whether the test compound would be likely to have that characteristic.

Example 5 – Use of an Experimentally Derived SDAR and ^{13}C NMR Spectral Data to Screen a Combinatorial Library of Compounds for Estrogen Receptor Binding Ability

Combinatorial chemistry refers to methods of generating large numbers of compounds from smaller building block compounds that play an important role in generating lead compounds for rational drug design. The building block compounds are allowed to react to yield new compounds, either by mixed

synthesis using all building blocks together at once or by sequential reaction of the building block compounds. The reactions can take place within the virtual environment of a computer or they can actually be carried out in a reactor system. The resulting set of compounds is termed a combinatorial library.

5 A combinatorial library of compounds is produced using a combinatorial chemistry software package such as Afferent Structure™ (Afferent Systems, San Francisco, CA). With this software package it is possible to begin with a molecule and perform virtual synthetic steps on that molecule and any intermediate compounds made therefrom to yield a large number of intermediate and final
10 product structures.

For each of the structures generated using the combinatorial chemistry software, ^{13}C NMR spectra are predicted. The ^{13}C NMR spectra may be predicted by any known method. Examples of methods for predicting ^{13}C NMR spectra include the neural network methods described by Kvasnicka (Kvasnicka, V., *J. Math. Chem.*, **6**: 63-76, 1991) and the quantum mechanical calculations of Dios et al. (Dios et al., *Science* **260**:1491-1496, 1993). Software for predicting ^{13}C NMR spectra is also available from Advanced Chemistry Development, Toronto, Ontario, Canada (www.acdlabs.com) (ACD/CNMR Spectrum Generator).
15

The predicted ^{13}C NMR spectra are segmented into bins as the experimental
20 ^{13}C NMR spectra in Example 1 were segmented. The segmented, predicted spectral data is input into the pattern recognition program and the SDAR established for estrogen receptor binding based on ^{13}C NMR only, from Example 1, is used to classify the compounds in the combinatorial library as either strong or medium estrogen receptor binders. Those compounds predicted by the SDAR to
25 be strong estrogen receptor binders may then be tested experimentally for ability to bind to estrogen receptors.

Combinatorial libraries offer a quick approach to generating large numbers of new compounds, yet screening those compounds for specific biological activities is difficult and time consuming. The methods of the present invention provide
30 rapid methods of screening the products produced by combinatorial chemistry

methods so that time consuming assays are performed only on compounds likely to exhibit the desired endpoint properties.

While this embodiment utilized an SDAR generated using experimental spectral data, in another embodiment, calculated spectral data is used to establish the SDAR. An SDAR established with calculated spectral data may be desirable when screening compounds according to their calculated spectral data, as the errors in calculated peak position are advantageously similar in both the training set spectral data and test compound spectral data.

10 **Example 6 - Screening of Fractions for Specific Activities using SDAR**

The SDAR methods of the present invention are useful for screening raw fractions of compounds derived either from natural sources or from chemical reaction mixtures (including experimental combinatorial libraries).

A biological source of potentially new compounds, such as a sponge, is homogenized and partitioned to provide an aqueous fraction and an organic fraction. Each of these fractions is then chromatographed by any known method to yield a larger number of fractions that contain one or more compounds. Within some of these fractions, a compound or compounds capable of binding to estrogen receptors may be present. SDAR is used to quickly screen the fractions for the presence of estrogen receptor binding compounds.

An unassigned ^{13}C NMR spectrum is obtained for each fraction and the spectrum is segmented into bins of 1ppm width. The fraction spectral data is then used along with the SDAR based on ^{13}C NMR data only, as in Example 1, to predict the fraction's ability to bind to estrogen receptors. Screening of fractions in this manner avoids the time-consuming and costly step of subjecting each fraction to a bioassay.

Those fractions identified as containing compounds having the potential to bind to estrogen receptors may then be subjected to further analysis to reveal the number and identity of the compounds within the fraction.

30

Example 7- Endpoints

Endpoints for use with the SDAR methods encompass the full range of biological, chemical, and physical properties exhibited by molecules. The methods of the present invention can be used to assist in drug design, biological activity
5 predictions, toxicological predictions, chemical reactivity predictions, and metabolic pathway predictions. An endpoint is any molecular property or activity that can be measured qualitatively or quantitatively. Endpoints may be expressed in absolute or relative terms.

Endpoints may be chosen to establish SDARs that can be used to predict
10 the environmental fate and toxicity of compounds. The ability of compounds to penetrate membranes, bind to enzyme active sites, react with soil, air, or water constituents, bind to soil constituents, hydrolyze, oxidize, and be transported in the environment can be used, along with spectral data for those compounds, to produce useful SDARs.

15 Spectral data can be used in combination with non-specific measures of toxicity, mutagenicity, teratogenicity, and carcinogenicity to establish SDARs. One example of such a non-specific measure is the Ames test. DNA damage and repair tests, Phosphorous-32 postlabeling, mutation induction in transgenes are others. Yet others include transgenic mouse assays, including the p53 +/- deficient
20 model, the Tg.AC model, the TgHras2 model, and the XPA deficient model. LD₅₀ and EC₅₀ may provide endpoints for SDAR methods as well. Alternatively, the ability of compounds to induce specific biological outcomes such as cellular changes can be chosen as the endpoint used to establish the SDAR. For example, relevant tissues may be examined for changes at the cellular level using
25 morphological, histochemical, or functional criteria. As appropriate, attention may be directed to such changes as the dose-relationships for apoptosis, cell proliferation, liver foci of cellular alteration, or changes in intercellular communication.

An SDAR may be established based upon any measurable response elicited
30 in animals, plants, and microbes upon exposure to a series of compounds. Examples include SDARs based upon antiviral and antimicrobial activity. The

ability of compounds to induce metabolic disorders such as alterations in sugar metabolism may provide a useful endpoint. Phytotoxicity and stimulation of plant growth and reproduction are other examples. Pesticidal activity is yet another example. Measures of anti-hypertensive activity, anti-pyretic activity, anti-depressant activity, and the like further illustrate useful endpoints that are usually related to human health.

Phototoxicity, both specific and non-specific, may be correlated with spectral features to yield an SDAR.

Multiple endpoints may be utilized to establish multiple SDARs from a single set of spectral data. Compounds then may be screened based upon their spectra using multiple SDARs for any combination of desirable or undesirable activities. One example of a useful combination is that of maximal potential efficacy as a therapeutic agent with minimal potential side effects. Agrochemicals may be screened using multiple SDARs for species-specific toxicities and tolerances.

An especially useful application of the methods of the present invention is to the prediction of ligand-target molecule binding. The binding of a molecule to a target such a protein, nucleic acid, synthetic polymer, chimeric molecule, or membrane constituent is often the most important step in the elicitation of a particular property or activity by a molecule. Binding affinities for ligand-target molecule interactions can be expressed in either absolute (e.g., an equilibrium constant) or in relative (e.g., relative to a reference compound, as determined for example by a competitive binding experiment) terms. Example 1 above is one example of how the relative binding affinity of a series of molecules can be utilized along with spectral data to establish a predictive SDAR model. SDAR models based upon relative binding affinities may be useful for rapidly and inexpensively screening compounds for a particular activity. They also may be useful tools for rational drug design when used to identify the spectral, and thus structural, features responsible for that activity.

The metabolic pathway involved in the production or destruction of a series of molecules is another endpoint useful for the methods of the present invention.

A predictive SDAR based upon pathway-structure relationships may be able to predict the biosynthetic path for newly discovered naturally occurring compounds. Similarly, SDAR using biodegradability as an endpoint may be useful for predicting the residence time of pollutants in the environment.

5 Rates of reaction and other measures of reactivity, such as site of reaction on a molecular structure, including the site of electrophilic aromatic substitutions on aromatic compounds, are useful chemical endpoints for the practice of the present invention.

Physical constants such as water-octanol partition coefficients, vapor
10 pressures, pKa, pKb, hydrophobicities, relative acidities and basicities as well as water solubilities can be used with spectral data to provide SDARs. Such estimates may be especially useful for physical properties that are difficult and time-consuming to measure. For example, octanol-water partition coefficients are important for modeling the environmental transport of chemicals. While the
15 octanol-water partition coefficient of a compound might be available, it is less likely that transient species derived from that compound during biodegradation are available in sufficient quantities to measure their octanol-water partition coefficients. SDAR according to the methods of the present invention provides an efficient way to predict the octanol-water partition coefficient for transient species,
20 whose environmental transport characteristics need to be modeled.

Other examples of endpoints useful for the methods of the present invention, in addition to those discussed above, may be found in Hansch and Leo, *Exploring QSAR: Fundamentals and Applications in Chemistry and Biology*, American Chemical Society, 1995. Further examples of endpoints useful for the
25 methods of the present invention may be found in *Quantitative Structure-Activity Relationships in Environmental Sciences-VII*, Chen and Schüürmann, eds., SETAC Press, 1997.

Example 8 – Spectral Data

30 Spectroscopy refers to branch of analytical chemistry in which atomic and molecular structure is studied by measuring radiant energy absorbed or emitted by

a substance in any of the wavelengths of the electromagnetic spectrum, in response to excitation by an external energy source. The types of absorption and emission spectroscopy are usually identified by the wavelength involved, such as gamma-ray, X-ray, UV, visible, infrared, microwave, and radiofrequency. Nuclear magnetic resonance spectroscopy (NMR) examines differences in energy states created by a magnetic field. Spectral data refers to the measurements of the energy differences across the spectrum, and spectral patterns refer to differences in the detected energy differences measured across a region of the electromagnetic spectrum. Any instrumental method that produces data that depend upon the structural and quantum mechanical properties of a molecule may be utilized with the methods of the present invention.

Spectral data as used in some embodiments includes the entire spectrum (or spectra) generated by the instrumental method (or methods) of spectroscopy or by calculation. Furthermore, the spectral data need not be assigned to particular structural features. In other embodiments the spectral data comprises only a portion of the spectrum or spectra available. The spectral portions utilized in the methods of the present invention may conveniently cover a spectral region known to typically arise from one or more particular structural features. For example, with respect to ^{13}C NMR spectral data, spectral data can be obtained from the entire ^{13}C NMR spectrum (0 to 220 ppm), or at least half or a third of that spectrum, or at least a 60 ppm, 80 ppm, 100, or 150 ppm portion of the spectrum. With respect to IR spectral data, for example, the spectral data can be obtained from the entire IR spectrum (4000 cm^{-1} to 500 cm^{-1}), or at least a hundredth, fiftieth, quarter, or half of that spectrum, or at least a 35, 50, 100, 200, 500, or 1000 cm^{-1} portion of the spectrum. In particular embodiments, all or substantially all of the spectral data within a particular portion of the spectrum is obtained. However, the spectral features within the entire spectrum or within the portion of the spectrum need not be assigned to structural features and referenced to the corresponding spectral features arising from the structure features of a reference compound. For example, rather than utilizing the assigned ^{13}C NMR resonances of the carbon atoms in a substituted benzene ring referenced to the ^{13}C NMR

resonances of the carbon atoms in unsubstituted benzene, the present invention could utilize the entire spectral region wherein benzene ring carbons typically fall, without assigning and referencing the data therein to the corresponding spectral data for benzene, which has a known structure.

5 Nuclear magnetic resonance (NMR) data is especially attractive for use with the present invention because of the large amount of structural information contained in a NMR spectrum. NMR instrumentation is widely available and NMR spectra are obtained routinely during structure elucidation. Additionally, the NMR spectra of many compounds have already been measured and are available
10 (see Example 1 for some representative sources).

¹³C NMR and ¹H NMR spectral data (especially ¹³C NMR spectral data) are very sensitive to subtle changes in substitution, conformation, chirality, and electronic density. Moreover, changes in ¹³C NMR chemical shifts can occur at a site as many as five carbon atoms removed from the site of the variation.
15 Solvation and proton-exchange effects on the electronic properties of molecules are more clearly reflected in ¹H NMR chemical shifts and line widths. One-dimensional ¹³C NMR and ¹H NMR spectral data as well as two-dimensional ¹³C-¹H heterocorrelation data such as that derived from HSQC and HMQC experiments are useful. Furthermore, modern FT-NMR instruments are capable
20 of providing NMR spectral data for as little as 1 nanogram of a compound (an amount likely insufficient for performing standard bioassays such as the Ames test).

 NMR data may be segmented into bins prior to their analysis, along with endpoint data, in a pattern-recognition program. Suitable bin widths will vary
25 according to the identity of the nuclei for which the spectrum is generated, and whether the technique is one or two-dimensional. For one-dimensional ¹³C NMR spectral data, the bin width may be varied from the digital resolution of the instrument (typically about 0.1 ppm) to about 50ppm. For one-dimensional ¹H NMR spectral data, the bin width may be varied from the instrumental digital
30 resolution (typically about 0.01 ppm) to about 2 ppm. For two-dimensional ¹³C-¹H heterocorrelation data, the bin may be defined by similar corresponding widths in

both the ^{13}C and ^1H dimensions. Even data of higher dimensions (e.g., three, four, etc.) including NMR spectral data from other nuclei, such as ^{15}N , ^{31}P , ^{19}F , ^{17}O , and ^{35}S may be used in establishing an SDAR. Correspondingly, bins may be defined with respect to each dimension and may be of a width equal to the digital
5 resolution of the data or greater.

In another embodiment of the invention, ^{13}C NMR spectral data are predicted by calculation (see, for example, Dios et al., *Science* **260**:1491-1496, 1993 and Kvasnicka, V., *J. Math. Chem.*, **6**: 63-76, 1991) and used in an SDAR model that has been trained on true ^{13}C NMR spectral data. Software for
10 predicting ^{13}C NMR spectra is also available from Advanced Chemistry Development, Toronto, Ontario, Canada (www.acdlabs.com) (ACD/CNMR Spectrum Generator). Predicted ^{13}C NMR spectral data may be used, for example, to aid in rational drug design by allowing proposed structures to be tested for potential activities before synthesis is attempted.

15 If predicted spectral data is utilized to establish an SDAR, the spectral data may be segmented into bins that are desirably of a width equal to the average standard deviation in chemical shift predicted by the method, or greater. Likewise, test structure predicted spectra may be segmented in a similar fashion.

Mass spectrometry can provide a measure of the size of a molecule, the
20 size and identity of a molecule's structural subunits, and information regarding bond strengths within a molecule. Mass spectral data, especially electron impact mass spectral (EI MS) data, has already been obtained for many compounds and, even more so than NMR data, is available from convenient sources (see Example 1). EI MS data is also a standard technique used in structure elucidation.

25 Other mass spectrometric techniques that are useful for providing additional and often complementary information include time-of-flight mass spectrometry (TOF MS), chemical ionization mass spectrometry (CI MS), fast-atom bombardment (FAB). Modern TOF MS spectrometers are capable of providing mass-spectral data from 1 ng or less of purified material (an amount that is likely
30 insufficient for performing standard activity screens such as the Ames test).

Mass spectral data may be segmented into ranges of m/z ratio (for instance, ranges corresponding to mass ranges from about the digital resolution of the instrumental method, typically about 0.1 amu, to about 50 amu) or may be segmented according to integer m/z ratios, with non-integer m/z ratios being
5 rounded to the nearest integer.

Infrared (IR) spectra may also be used, for example in establishing SDAR models capable of discerning differences in activity seen for tautomers that are indistinguishable in NMR and mass spectrometric data. Infrared (IR) spectra are treated in a similar fashion to the NMR spectral data of Example 1 in that each
10 spectrum may be separated into bins of a certain spectral range, for example from about 1 cm^{-1} to about 200 cm^{-1} for entry into a pattern-recognition program

Ultraviolet-Visible (UV-Vis) spectral data, which is inherently reflective of the electronic energy levels of a molecule, may be used by segmenting the spectral data into bins of a certain spectral range, for example from about the digital
15 resolution of the instrumental method to about 50 nm. UV-Vis spectral data can be used, for example, in predicting phototoxicity under solar illumination. Similarly, fluorescence and phosphorescence spectra may be handled analogously to UV-Vis spectra and utilized to establish an SDAR. Fluorescence and phosphorescence spectra reflect the energy redistribution within a molecule upon
20 absorption of light and thus may provide important structure descriptors for predicting the light driven properties of molecules.

In addition to the possibility of using a single type of spectral data to establish an SDAR, spectral data of various types may be combined to form composite sets of spectral data. Entire spectra or particular regions of spectra may
25 be combined to yield spectral data sets that may be used in the methods of the present invention, along with endpoint data, to establish the SDAR. Spectral data may come from any composite of NMR, MS, IR, Fluorescence, Phosphorescence, and UV-Vis spectra, including composites of different species of spectra within these broad genera of spectra. Furthermore, different regions of any type of
30 spectrum can be segmented into bins of different sizes so, for example, portions of a spectrum with many closely spaced peaks can be described by narrow spectral

bins and portions of a spectrum without many peaks can be described by wide spectral bins.

In some embodiments the spectral data is not used in its raw form to establish an SDAR, but rather the data is subjected to pattern recognition analysis after some sort of pre-treatment to improve the ability of pattern recognition to extract the SDAR. For example normalization may be used to equalize the importance of spectral data derived from different instrumental methods when forming a composite, such as, a composite of MS data and NMR data wherein the maximum signals might be 100 and 1000 respectively. Autoscaling may be used to equalize the importance of inherently weak spectral data with inherently strong spectral data, for example, UV-Vis absorption bands within an absorption spectrum with very different extinction coefficients. Fisher-weighting may be used to emphasize the spectral data that are most important for predicting the endpoint data, such as spectral data found in compounds with a large endpoint values but absent from compounds with small endpoint values.

In general, techniques for pre-treating data include artifact removal and/or linearization, centering, and scaling and weighting. A common form of artifact removal is baseline correction of a spectrum. Common linearizations include the conversion of spectral transmittance into spectral absorbance and the multiplicative scatter correction for diffuse reflectance spectra. Centering, sometimes called mean centering is simply the subtraction of the mean spectral signal at each frequency or m/z from each spectrum. Scaling or weighting involves multiplying all of the spectra by a different scaling factor for each sub-spectral region. This is done to increase or decrease the influence of certain spectral regions or features. A particular example of weighting is Fisher-weighting. Two types of scaling are typically encountered, variance scaling and autoscaling. Further discussion of data pretreatment may be found in Kramer, R., *Chemometric Techniques for Quantitative Analysis*, Marcel Dekker, Inc., 1998. Additional methods for the pre-treatment of data prior to pattern recognition are known in the art and are included in the present invention.

Since spectral data is utilized in some embodiments in the same manner as 3D-QSAR utilizes comprehensive descriptors for structural and statistical analyses (CODESSA) (see, Tong et al., *J. Med. Chem.*, **39**: 380-387, 1995 and Collantes et al., *J. Anal. Chem.*, **68**: 2038-2043, 1996, both of which are incorporated herein
5 by reference) it is possible to combine the spectrally derived structure descriptors of the present invention with structure descriptors based on bulk measurements (such as the octanol/water partition coefficient) and still retain the ability to produce predictive models without the need for structural knowledge beforehand. Hybrid methods also fall within the scope of the invention wherein the spectrally
10 derived descriptors, which do not require structural knowledge beforehand, may be combined with other structure descriptors that require structural knowledge to produce a larger set of descriptors for use in a predictive model.

Example 9 – Pattern-Recognition Programs

15 Pattern-recognition programs useful for practicing the present invention are of two major types; statistical and artificial intelligence.

Statistical methods include Principal Component Analysis (PCA) and variations of PCA such as linear regression analysis, cluster analysis, canonical variates, and discriminant analysis, soft independent models of class analogy
20 (SIMCA), expert systems, and auto spin (see, for example, Harrington, *RESolve Software Manual*, Colorado School of Mines, 1988, incorporated by reference). Other examples of statistical analysis software available for principal-component-based methods include SPSS (SPSS Inc., Chicago, IL), JMP (SAS Inc., Cary NC), Stata (Stata Inc., College Station, TX) and Cluster (available to run from
25 entropy:~dblank/public_html/cluster).

Artificial intelligence methods include neural networks and fuzzy logic. Neural networks may be one-layer or multilayer in architecture (See, for example, Zupan and Gasteiger, *Neural Networks for Chemists*, VCH, 1993, incorporated herein by reference). Examples of one-layer networks include Hopfield networks,
30 Adaptive Bidirectional Associative Memory (ABAM), and Kohonen Networks. Examples of Multilayer Networks include those that learn by counter-propagation

and back-propagation of error. Artificial neural network software is available from, among other sources, Neurodimension, Inc., Gainesville, FL (Neurosolutions) and The Mathworks, Inc., Natick, MA (MATLAB Neural Network Toolbox).

5 Spectral patterns can be analyzed using other approaches. Analog spectral peak patterns may be digitized, and image analysis may be used to search for similarities or differences between the spectral patterns of training sets and test compounds

10 **Example 10 – Computer Environments for SDAR Methods.**

The SDAR methods of the present invention may be implemented using a single computer or utilizing a distributed computing environment.

Exemplary Distributed Computing Environment

Fig. 13 illustrates a distributed computing environment in which the software elements used to implement the SDAR methods of the present invention may reside. The distributed computing environment 100 includes two computer systems 102, 104 connected by a connection medium 106. The computer systems 102, 104 can be any of several types of computer system configurations, including personal computers, multiprocessor systems, and the like. In terms of logical relation with other computer systems, a computer system can be a client, a server, a router, a peer device, or other common network node. Moreover, although Fig. 13 illustrates two computer systems 102, 104, the present invention is equally applicable to an arbitrary, larger number of computer systems connected by the connection medium 106. Additional computer systems 102 or 104 may be connected by an arbitrary number of connection mediums 106. The connection medium 106 can comprise any local area network (LAN), wide area network (WAN), or other computer network, including but not limited to Ethernets, enterprise-wide computer networks, intranets and the Internet.

Portions of the SDAR software can be implemented in a single computer system 102 or 104, with the application later distributed to other computer systems 102, 104 in the distributed computing environment 100. Portions of the SDAR

software may also be practiced in a distributed computing environment 100 where tasks are performed by a single computer system 102 or 104 acting as a remote processing device that is accessed through a communications network, with the distributed application later distributed to other computer systems in the distributed computing environment 100. In a networked environment, program modules comprising the SDAR software can be located on more than one computer system 102 or 104. Communication between the computer systems in the distributed computing network may advantageously include encryption of the communicated data.

10

Exemplary Computer System

Fig. 10 illustrates an example of a computer system 120 that can serve as an operating environment for the SDAR software. With reference to Fig. 14 an exemplary computer system for implementing the invention includes a computer 120 (such as a personal computer, laptop, palmtop, set-top, server, mainframe, and other varieties of computer), including a processing unit 121, a system memory 122, and a system bus 123 that couples various system components including the system memory to the processing unit 121. The processing unit can be any of various commercially available processors, including Intel x86, Pentium and compatible microprocessors from Intel and others, including Cyrix, AMD and Nexgen; Alpha from Digital; MIPS from MIPS Technology, NEC, IDT, Siemens, and others; and the PowerPC from IBM and Motorola. Dual microprocessors and other multi-processor architectures also can be used as the processing unit 121.

The system bus can be any of several types of bus structure including a memory bus or memory controller, a peripheral bus, and a local bus using any of a variety of conventional bus architectures such as PCI, VESA, AGP, Microchannel, ISA and EISA, to name a few. The system memory includes read only memory (ROM) 124 and random access memory (RAM) 125. A basic input/output system (BIOS), containing the basic routines that help to transfer information between elements within the computer 120, such as during start-up, is stored in ROM 124.

The computer 120 further includes a hard disk drive 127, a magnetic disk drive 128, e.g., to read from or write to a removable disk 129, and an optical disk drive 130, e.g., for reading a CD-ROM disk 131 or to read from or write to other optical media. The hard disk drive 127, magnetic disk drive 128, and optical disk drive 130 are connected to the system bus 123 by a hard disk drive interface 132, a magnetic disk drive interface 133, and an optical drive interface 134, respectively. The drives and their associated computer-readable media provide nonvolatile storage of data, data structures, computer-executable instructions, etc. for the computer 120. Although the description of computer-readable media above refers to a hard disk, a removable magnetic disk and a CD, it should be appreciated by those skilled in the art that other types of media which are readable by a computer, such as magnetic cassettes, flash memory cards, digital video disks, Bernoulli cartridges, and the like, can also be used in the exemplary operating environment.

A number of the SDAR program modules can be stored in the drives and RAM 125, including an operating system 135, one or more application programs 136, other program modules 137, and program data 138.

A user can enter commands and information into the computer 120 through a keyboard 140 and pointing device, such as a mouse 142. Other input devices (not shown) can include a microphone, satellite dish, scanner, or the like. These and other input devices are often connected to the processing unit 121 through a serial port interface 146 that is coupled to the system bus, but can be connected by other interfaces, such as a parallel port, game port or a universal serial bus (USB). A monitor 147 or other type of display device is also connected to the system bus 123 via an interface, such as a video adapter 148. In addition to the monitor, computers typically include other peripheral output devices (not shown), such as printers.

The computer 120 can operate in a networked environment using logical connections to one or more other computer systems, such as computer 102. The other computer systems can be servers, routers, peer devices or other common network nodes, and typically include many or all of the elements described relative to the computer 120, although only a memory storage device 149 has been

illustrated in Fig. 14. The logical connections depicted in Fig. 10 include a local area network (LAN) 151 and a wide area network (WAN) 152. Such networking environments are common in offices, enterprise-wide computer networks, intranets and the Internet.

5 When used in a LAN networking environment, the computer 120 is connected to the local network 151 through a network interface or adapter 153. When used in a WAN networking environment, the computer 120 typically includes a modem 154 or other means for establishing communications (e.g., via the LAN 151 and a gateway or proxy server 155) over the wide area network 152,
10 such as the Internet. The modem 154, which can be internal or external, is connected to the system bus 123 via the serial port interface 146. In a networked environment, program modules depicted relative to the computer 120, or portions thereof, can be stored in the remote memory storage device. It will be appreciated that the network connections shown are exemplary and other means of establishing
15 a communications link between the computer systems (including an Ethernet card, ISDN terminal adapter, ADSL modem, 10BaseT adapter, 100BaseT adapter, ATM adapter, or the like) can be used.

 In accordance with the practices of persons skilled in the art of computer programming, a particular embodiment of the SDAR method is described in Fig. 4
20 with reference to acts and symbolic representations of operations that may be performed by the computer 120. Such acts and operations are sometimes referred to as being computer-executed. It will be appreciated that the acts and symbolically represented operations include the manipulation by the processing unit 121 of electrical signals representing data bits which causes a resulting
25 transformation or reduction of the electrical signal representation, and the maintenance of data bits at memory locations in the memory system (including the system memory 122, hard drive 127, floppy disks 129, and CD-ROM 131) to thereby reconfigure or otherwise alter the computer system's operation, as well as other processing of signals. The memory locations where data bits are maintained
30 are physical locations that have particular electrical, magnetic, or optical properties corresponding to the data bits.

In view of the many possible embodiments to which the principles of our invention may be applied, it should be recognized that the illustrated embodiments are only specific examples of the invention and should not be taken as a limitation on the scope of the invention. Rather, the scope of the invention is defined by the
5 following claims. We therefore claim as our invention all that comes within the scope and spirit of these claims.

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207

We Claim:

1. A method of predicting a biological activity of a molecule,
comprising:
 - 5 obtaining spectral data for a test compound;
comparing the spectral data of the test compound to a pattern of spectral
data associated with a biological activity, derived not exclusively from the assigned
spectral data of a training set of compounds having a known biological activity;
detecting similarities between the pattern of spectral data associated with a
10 biological activity of the training set and a pattern of spectral data for the test
compound to determine whether the test compound is predicted to share the
biological activity.
 2. The method of claim 1 wherein the spectral data are obtained
without first correlating the spectral data with corresponding structural features.
 - 15 3. The method of claim 1 wherein the pattern of spectral data
associated with a biological activity is derived without first correlating the spectral
data with corresponding structural features.
 4. The method of claim 1, wherein the pattern of spectral data of the
training set is a pattern obtained by separating the spectral data of the training set
20 of compounds into sub-spectral units.
 5. The method of claim 4, wherein the pattern of spectral data of the
test compound is obtained by separating the spectral data of the test compound into
substantially the same sub-spectral units into which the spectral data of the training
set is separated.
 - 25 6. The method of claim 1, wherein the spectral data is one type of
spectral data.
 7. The method of claim 6, wherein the spectral data comprises one of
nuclear magnetic resonance, mass spectral, infrared, ultraviolet-visible,
fluorescence, or phosphorescence data.
 - 30 8. The method of claim 1, wherein the spectral data is a composite of
different types of spectral data.

9. The method of claim 8, wherein the different types of spectral data comprise two or more of the group consisting of nuclear magnetic spectroscopy (NMR), mass spectroscopy (MS), infrared (IR) spectroscopy, and ultraviolet-visible (UV-Vis) spectroscopy.

5 10. The method of claim 1, wherein the spectral pattern of the test compound and the spectral pattern of the training set are segmented into sub-spectral units, and the spectral data of the training set is scaled to normalize the importance of different signals within the spectral data of the training set prior to deriving a pattern associated with a biological activity.

10 11. The method of claim 10, wherein the scaling is auto-scaling.

12. The method of claim 10, wherein the spectral data of the training set is weighted to emphasize signals that are important for determining the endpoint class of compounds in the training set before deriving a pattern associated with a biological activity.

15 13. The method of claim 12 wherein the weighting is Fisher-weighting.

14. The method of claim 1, wherein detecting similarities between the pattern of spectral data associated with a biological activity of the training set and the pattern of spectral data for the test compound comprises performing computer implemented pattern recognition.

20 15. The method of claim 1, wherein detecting similarities between the pattern of spectral data associated with a biological activity of the training set and the pattern of spectral data for the test compound comprises detecting relative intensities of signals associated with one or more of the sub-units of the spectrum of the training set, and detecting relative intensities of signals associated with the
25 same one or more sub-units of a spectrum of the test compound.

16. The method of claim 15, wherein the relative intensities are canonical variate factors of the spectral data associated with a biological activity of the training set and the spectral signals of the test compound.

17. The method of claim 1, wherein the method is computer
30 implemented.

18. A computer implemented system for predicting a biological activity of a test compound, comprising:

receiving as input spectral data for a test compound;

receiving as input spectral data of a training set of compounds having a
5 known biological activity; and

comparing the pattern of spectral data of the training set associated with the biological activity to the spectral pattern of the test compound to determine whether the spectral pattern of the test compound matches the spectral pattern associated with the biological activity of the training set.

10 19. The computer implemented system of claim 18, wherein comparing the spectral patterns comprises comparing the spectral patterns with computer implemented pattern recognition programs.

20. The computer implemented system of claim 19, wherein the spectral data for the test compound and the spectral data for the training set are divided into
15 substantially identical spectral bins, so that a signal within individual spectral bins is compared between spectral patterns of the training set associated with the biological activity and the test compound.

21. The computer implemented system of claim 18, wherein the spectral patterns are obtained by inputting spectral data selected from the group consisting
20 of nuclear magnetic resonance data, mass spectral data, infrared data, ultraviolet-visible data, fluorescence data, phosphorescence data, and composites of two or more such spectral data.

22. The computer implemented system of claim 21, wherein the spectral data for the training set are converted into canonical variates associated with the
25 biological activity of the training set, and the spectral data for the test compound are compared to the canonical variates of the training set spectral data.

23. The computer implemented system of claim 22, wherein the biological activity is binding affinity to a hormone receptor, and the canonical variates for the training set include peaks in bins that are associated with hormone
30 receptor binding of a pre-selected affinity.

24. The computer implemented system of claim 23, wherein the spectral data comprise nuclear magnetic resonance data and mass spectral data.

25. A computer readable medium having stored thereon instructions for performing the actions of claim 1.

5 26. A computer readable medium having stored thereon instructions for performing the actions of claim 18.

27. A method for predicting a biological, chemical, or physical property of a molecule, comprising:

providing spectral data segmented into spectral sub-units, for a plurality of
10 training compounds;

inputting the segmented spectral data and endpoint data into a pattern-recognition program;

training the pattern-recognition program with the segmented spectral data and endpoint data to establish a relationship between the spectral sub-units of the
15 segmented spectral data and the endpoint;

providing segmented spectral data for a test compound that is segmented into substantially the same spectral sub-units as the spectral data of the training compounds; and

comparing the relationship between the spectral subunits of the segmented
20 spectral data and the endpoint to the spectral subunits of the test compound's segmented spectral data to predict the endpoint of the test compound.

28. The method of claim 27, wherein knowledge of the structures of the training compounds and the test compound are not necessarily known beforehand.

29. The method of claim 27 wherein the segmented spectral data of the
25 training set is autoscaled and Fisher-weighted before training the pattern recognition program.

30. The method of claim 27 wherein the spectral data is chosen from the group consisting of nuclear magnetic resonance data, mass spectral data, infrared data, UV-Vis data, fluorescence data, phosphorescence data, and composites
30 thereof.

31. The method of claim 30 wherein the spectral data is chosen from the group consisting of ^{13}C NMR data, EI MS data, and composites thereof.

32. The method of claim 27, wherein the endpoint is a ligand-target molecule-binding affinity.

5 33. The method of claim 32, wherein the ligand-target molecule binding affinity is an estrogen-receptor binding affinity.

34. The method of claim 27, wherein the endpoint is selected from the group consisting of:

a measure of biodegradability;

10 a measure of toxicity;

participation in a metabolic pathway;

a partition coefficient;

a reaction rate;

a quantum yield;

15 a measure of phototoxicity;

an equilibrium constant; and

a site of reaction on a molecular structure.

35. The method of claim 34, wherein the partition coefficient is the octanol-water partition coefficient.

20 36. The method of claim 27, wherein non-spectral structure descriptors that do not necessarily depend upon structural knowledge beforehand are provided for the test compound and the training compounds; used to establish a relationship between the segmented spectral data, the non-spectral structure descriptors, and the endpoint for the training compounds; and used to predict the endpoint of the
25 test compound.

37. The method of claim 36, wherein the non-spectral structure descriptors are chosen from the group consisting of partition coefficients, solubilities, relative acidities, relative basicities, pKa, pKb, reaction rates, and equilibrium constants.

30 38. The method of claim 37, wherein the partition coefficient is the octanol-water partition coefficient.

39. The method of claim 27, wherein non-spectral structure descriptors that do depend upon structural knowledge beforehand are provided for the test compound and the training compounds; used to establish a relationship between the segmented spectral data, the non-spectral structure descriptors, and the endpoint
5 for the training compounds; and used to predict the endpoint of the test compound.

40. The method of claim 39, wherein the non-spectral descriptors are calculated using a quantum mechanical or electrostatic potential method.

41. A method for using spectral data as a set of structure descriptors for a compound that does not necessarily require knowledge of the compound's structure
10 beforehand, comprising:

providing spectral data; and
segmenting the spectral data into bins.

42. A method for establishing a relationship between spectral data and a biological, chemical, or physical property, comprising:
15 providing spectral data;
segmenting the spectral data into bins.
detecting patterns in the bins of the spectral data that are associated with the property.

43. The method of claim 42, further comprising detecting corresponding
20 patterns in spectral data of test compounds to select test compounds having the property.

44. The method of claim 43, wherein the test compounds are mixtures of compounds.

45. The method of claim 42 including one or more of: auto-scaling the
25 segmented data; and weighting the segmented spectral data.

46. The method of claim 45 wherein weighting of the segmented data comprises Fisher-weighting of the segmented spectral data.

47. The method of claim 1, wherein the biological activity of the test compound is predicted without reference to a chemical structure of the test
30 compound.

48. A method for establishing a spectral data activity relationship, comprising;

providing endpoint data for a plurality of compounds;

providing spectral data for a plurality of compounds;

5 segmenting the spectral data for the plurality of compounds into bins;

autoscaling the numerical data obtained from the spectral features within each of the bins;

Fisher-weighting the data within each of the bins; and

correlating information in the bins and the endpoint using a means for

10 pattern recognition.

49. The method of claim 47 wherein the spectral data is selected from the group consisting of NMR data, MS data, IR data, UV-Vis data, fluorescence data, phosphorescence data, and composites thereof.

50. The method of claim 48 wherein two or more types of spectral data

15 are normalized to each other in a composite.

51. A method for determining the structural features of a plurality of compounds that contribute to determining a particular endpoint property exhibited by the compounds, comprising:

providing segmented spectral data for the plurality of compounds;

20 providing endpoint data for the plurality of compounds;

establishing a spectral data-activity relationship by identifying the segmented spectral features that bias toward increased endpoint values and the segmented spectral features that bias toward decreased endpoint values; and

identifying the structural feature leading to the segmented spectral features

25 that bias toward increased or decreased endpoint values for the plurality of compounds.

52. The method of claim 51 wherein the segmented spectral data is selected from the group consisting of NMR data, MS data, IR data, UV-Vis data, fluorescence data, phosphorescence data, and composites thereof.

53. The method of claim 52 wherein two or more types of segmented

30 spectral data are normalized to each other in a composite.

54. The method of claim 52 wherein the segmented spectral data is a composite of ^{13}C NMR and EI MS data.

0123456789101112131415161718192021222324252627282930313233343536373839404142434445464748495051525354555657585960616263646566676869707172737475767778798081828384858687888990919293949596979899100101102103104105106107108109110111112113114115116117118119120121122123124125126127128129130131132133134135136137138139140141142143144145146147148149150151152153154155156157158159160161162163164165166167168169170171172173174175176177178179180181182183184185186187188189190191192193194195196197198199200201202203204205206207208209210211212213214215216217218219220221222223224225226227228229230231232233234235236237238239240241242243244245246247248249250251252253254255256257258259260261262263264265266267268269270271272273274275276277278279280281282283284285286287288289290291292293294295296297298299300301302303304305306307308309310311312313314315316317318319320321322323324325326327328329330331332333334335336337338339340341342343344345346347348349350351352353354355356357358359360361362363364365366367368369370371372373374375376377378379380381382383384385386387388389390391392393394395396397398399400401402403404405406407408409410411412413414415416417418419420421422423424425426427428429430431432433434435436437438439440441442443444445446447448449450451452453454455456457458459460461462463464465466467468469470471472473474475476477478479480481482483484485486487488489490491492493494495496497498499500501502503504505506507508509510511512513514515516517518519520521522523524525526527528529530531532533534535536537538539540541542543544545546547548549550551552553554555556557558559560561562563564565566567568569570571572573574575576577578579580581582583584585586587588589590591592593594595596597598599600601602603604605606607608609610611612613614615616617618619620621622623624625626627628629630631632633634635636637638639640641642643644645646647648649650651652653654655656657658659660661662663664665666667668669670671672673674675676677678679680681682683684685686687688689690691692693694695696697698699700701702703704705706707708709710711712713714715716717718719720721722723724725726727728729730731732733734735736737738739740741742743744745746747748749750751752753754755756757758759760761762763764765766767768769770771772773774775776777778779780781782783784785786787788789790791792793794795796797798799800801802803804805806807808809810811812813814815816817818819820821822823824825826827828829830831832833834835836837838839840841842843844845846847848849850851852853854855856857858859860861862863864865866867868869870871872873874875876877878879880881882883884885886887888889890891892893894895896897898899900901902903904905906907908909910911912913914915916917918919920921922923924925926927928929930931932933934935936937938939940941942943944945946947948949950951952953954955956957958959960961962963964965966967968969970971972973974975976977978979980981982983984985986987988989990991992993994995996997998999100010011002100310041005100610071008100910101011101210131014101510161017101810191020102110221023102410251026102710281029103010311032103310341035103610371038103910401041104210431044104510461047104810491050105110521053105410551056105710581059106010611062106310641065106610671068106910701071107210731074107510761077107810791080108110821083108410851086108710881089109010911092109310941095109610971098109911001101110211031104110511061107110811091110111111121113111411151116111711181119112011211122112311241125112611271128112911301131113211331134113511361137113811391140114111421143114411451146114711481149115011511152115311541155115611571158115911601161116211631164116511661167116811691170117111721173117411751176117711781179118011811182118311841185118611871188118911901191119211931194119511961197119811991200120112021203120412051206120712081209121012111212121312141215121612171218121912201221122212231224122512261227122812291230123112321233123412351236123712381239124012411242124312441245124612471248124912501251125212531254125512561257125812591260126112621263126412651266126712681269127012711272127312741275127612771278127912801281128212831284128512861287128812891290129112921293129412951296129712981299130013011302130313041305130613071308130913101311131213131314131513161317131813191320132113221323132413251326132713281329133013311332133313341335133613371338133913401341134213431344134513461347134813491350135113521353135413551356135713581359136013611362136313641365136613671368136913701371137213731374137513761377137813791380138113821383138413851386138713881389139013911392139313941395139613971398139914001401140214031404140514061407140814091410141114121413141414151416141714181419142014211422142314241425142614271428142914301431143214331434143514361437143814391440144114421443144414451446144714481449145014511452145314541455145614571458145914601461146214631464146514661467146814691470147114721473147414751476147714781479148014811482148314841485148614871488148914901491149214931494149514961497149814991500150115021503150415051506150715081509151015111512151315141515151615171518151915201521152215231524152515261527152815291530153115321533153415351536153715381539154015411542154315441545154615471548154915501551155215531554155515561557155815591560156115621563156415651566156715681569157015711572157315741575157615771578157915801581158215831584158515861587158815891590159115921593159415951596159715981599160016011602160316041605160616071608160916101611161216131614161516161617161816191620162116221623162416251626162716281629163016311632163316341635163616371638163916401641164216431644164516461647164816491650165116521653165416551656165716581659166016611662166316641665166616671668166916701671167216731674167516761677167816791680168116821683168416851686168716881689169016911692169316941695169616971698169917001701170217031704170517061707170817091710171117121713171417151716171717181719172017211722172317241725172617271728172917301731173217331734173517361737173817391740174117421743174417451746174717481749175017511752175317541755175617571758175917601761176217631764176517661767176817691770177117721773177417751776177717781779178017811782178317841785178617871788178917901791179217931794179517961797179817991800180118021803180418051806180718081809181018111812181318141815181618171818181918201821182218231824182518261827182818291830183118321833183418351836183718381839184018411842184318441845184618471848184918501851185218531854185518561857185818591860186118621863186418651866186718681869187018711872187318741875187618771878187918801881188218831884188518861887188818891890189118921893189418951896189718981899190019011902190319041905190619071908190919101911191219131914191519161917191819191920192119221923192419251926192719281929193019311932193319341935193619371938193919401941194219431944194519461947194819491950195119521953195419551956195719581959196019611962196319641965196619671968196919701971197219731974197519761977197819791980198119821983198419851986198719881989199019911992199319941995199619971998199920002001200220032004200520062007200820092010201120122013201420152016201720182019202020212022202320242025202620272028202920302031203220332034203520362037203820392040204120422043204420452046204720482049205020512052205320542055205620572058205920602061206220632064206520662067206820692070207120722073207420752076207720782079208020812082208320842085208620872088208920902091209220932094209520962097209820992100210121022103210421052106210721082109211021112112211321142115211621172118211921202121212221232124212521262127212821292130213121322133213421352136213721382139214021412142214321442145214621472148214921502151215221532154215521562157215821592160216121622163216421652166216721682169217021712172217321742175217621772178217921802181218221832184218521862187218821892190219121922193219421952196219721982199220022012202220322042205220622072208220922102211221222132214221522162217221822192220222122222223222422252226222722282229223022312232223322342235223622372238223922402241224222432244224522462247224822492250225122522253225422552256225722582259226022612262226322642265226622672268226922702271227222732274227522762277227822792280228122822283228422852286228722882289229022912292229322942295229622972298229923002301230223032304230523062307230823092310231123122313231423152316231723182319232023212322232323242325232623272328232923302331233223332334233523362337233823392340234123422343234423452346234723482349235023512352235323542355235623572358235923602361236223632364236523662367236823692370237123722373237423752376237723782379238023812382238323842385238623872388238923902391239223932394239523962397239823992400240124022403240424052406240724082409241024112412241324142415241624172418241924202421242224232424242524262427242824292430243124322433243424352436243724382439244024412442244324442445244624472448244924502451245224532454245524562457245824592460246124622463246424652466246724682469247024712472247324742475247624772478247924802481248224832484248524862487248824892490249124922493249424952496249724982499250025012502250325042505250625072508250925102511251225132514251525162517251825192520252125222523252425252526252725282529253025312532253325342535253625372538253925402541254225432544254525462547254825492550255125522553255425552556255725582559256025612562256325642565256625672568256925702571257225732574257525762577257825792580258125822583258425852586258725882589259025912592259325942595259625972598259926002601260226032604260526062607260826092610261126122613261426152616261726182619262026212622262326242625262626272628262926302631263226332634263526362637263826392640264126422643264426452646264726482649265026512652265326542655265626572658265926602661266226632664266526662667266826692670267126722673267426752676267726782679268026812682268326842685268626872688268926902691269226932694269526962697269826992700270127022703270427052706270727082709271027112712271327142715271627172718271927202721272227232724272527262727272827292730273127322733273427352736273727382739274027412742274327442745274627472748274927502751275227532754275527562757275827592760276127622763276427652766276727682769277027712772277327742775277627772778277927802781278227832784278527862787278827892790279127922793279427952796279727982799280028012802280328042805280628072808280928102811281228132814281528162817281828192820282128222823282428252826282728282829283028312832283328342835283628372838283928402841284228432844284528462847284828492850285128522853285428552856285728582859286028612862286328642865286628672868286928702871287228732874287528762877287828792880288128822883288428852886288728882889289028912892289328942895289628972898289929002901290229032904290529062907290829092910291129122913291429152916291729182919292029212922292329242925292629272928292929302931293229332934293529362937293829392940294129422943294429452946294729482949295029512952295329542955295629572958295929602961296229632964296529662967296829692970297129722973297429752976297729782979298029812982298329842985298629872988298929902991299229932994299529962997299829993000300130023003300430053006300730083009301030113012301330143015301630173018301930203021302230233024302530263027302830293030303130323033303430353036303730383039304030413042304330443045304630473048304930503051305230533054305530563057305830593060306130623063306430653066306730683069307030713072307330743075307630773078307930803081308230833084308530863087308830893090309130923093309430953096309730983099310031013102310331043105310631073108310931103111311231133114311531163117311831193120312131223123312431253126312731283129313031313132313331343135313631373138313931403141314231433144314531463147314831493150315131523153315431553156315731583159316031613162316331643165316631673168316931703171317231733174317531763177317831793180318131823183318431853186318731883189319031913192319331943195319631973198319932003201320232033204320532063207320832093210321132123213321432153216321732183219322032213222322332234322532263227322832293230323132323233323432353236323732383239324032413242324332443245324632473248324932503251325232533254325532563257325832593260326132623263326432653266326732683269327032713272327332743275327632773278327932803281328232833284328532863287328832893290329132923293329432953296329732983299330033013302330333043305330633073308330933103311331

Abstract

Methods are disclosed for establishing a quantitative relationship between spectral properties of molecules and a biological, chemical, or physical endpoint of the molecules. Spectral data including data from nuclear magnetic resonance, mass spectrometric, infrared, and ultraviolet-visible techniques are used along with endpoint data to train a pattern-recognition program. The training yields a spectral data-activity relationship that may be used to predict the endpoint value of a molecule from its spectral data alone. Methods for rapidly screening isolated compounds or mixtures of compounds based upon their spectral data are included.

SDBS-¹³C NMR SDBS No. 1716 CDS-12-007
 C₁₃ H₁₆ O₂
 4,4'-isopropylidenediphenol

22.53 MHz
 0.025 g 0.5 ml DMSO-d₆





Fig. 1 (a)

SDBS-Mass

MS-NW-4834 SDBS NO. 1716
 4,4'-ISOPROPYLIDENEDIPHENOL
 C₁₃H₁₆O₂ (Mass of Molecular Ion 228)



Source Temperature 190 °C
 Sample Temperature 140 °C
 Direct 75 eV

Fig. 1 (b)

HIT-NO-1641 SCORE= () SDBS-NO-1716 [IR-N10A-05052 : KBR DISC]
 4,4'-ISOPROPYLIDENEDIPHENOL
 C₁₃H₁₆O₂





Fig. 1 (c)


¹³C

| δ ppm | Int. | Assign. |
|--------|------|---------|
| 154.72 | 411 | 1 |
| 140.94 | 330 | 2 |
| 127.11 | 1000 | 3 |
| 114.43 | 977 | 4 |
| 40.78 | 223 | 5 |
| 30.73 | 328 | 6 |

Fig 2(a)

SDBS-Mass

MS-NW-6834 SDBS NO. ~ 1716
 4,4'-ISOPROPYLIDENEDIPHENOL
 C15H16O2 (Mass of molecular ion: 228)

m/z ratio

| | |
|-------|----|
| 27.0 | 1 |
| 39.0 | 4 |
| 41.0 | 3 |
| 51.0 | 2 |
| 55.0 | 2 |
| 63.0 | 2 |
| 65.0 | 5 |
| 66.0 | 1 |
| 76.0 | 1 |
| 77.0 | 3 |
| 79.0 | 1 |
| 89.0 | 1 |
| 90.0 | 1 |
| 91.0 | 9 |
| 92.0 | 1 |
| 94.0 | 2 |
| 95.0 | 1 |
| 99.0 | 4 |
| 105.0 | 1 |
| 106.5 | 2 |
| 107.0 | 4 |
| 114.0 | 1 |
| 115.0 | 1 |
| 119.0 | 13 |
| 120.0 | 2 |
| 134.0 | 1 |
| 135.0 | 4 |
| 152.0 | 1 |
| 165.0 | 2 |
| 169.0 | 1 |
| 181.0 | 1 |
| 183.0 | 1 |
| 195.0 | 1 |
| 197.0 | 1 |
| 198.0 | 1 |
| 212.0 | 1 |
| 213.0 | 10 |
| 214.0 | 10 |
| 215.0 | 1 |
| 228.0 | 33 |
| 229.0 | 5 |

Fig 2(b)

| cm ⁻¹ | %T | cm ⁻¹ | %T | cm ⁻¹ | %T | cm ⁻¹ | %T | cm ⁻¹ | %T |
|------------------|----|------------------|----|------------------|----|------------------|----|------------------|----|
| 3168 | 21 | 2933 | 72 | 1436 | 49 | 1178 | 20 | 815 | 66 |
| 3070 | 74 | 2871 | 77 | 1384 | 62 | 1150 | 74 | 753 | 60 |
| 3050 | 74 | 1612 | 39 | 1363 | 47 | 1113 | 72 | 735 | 77 |
| 3030 | 70 | 1600 | 37 | 1296 | 66 | 1102 | 72 | 724 | 72 |
| 2975 | 46 | 1510 | 4 | 1247 | 14 | 1085 | 57 | 650 | 58 |
| 2966 | 46 | 1463 | 62 | 1239 | 12 | 1013 | 62 | 565 | 41 |
| 2956 | 62 | 1447 | 42 | 1221 | 19 | 827 | 14 | 663 | 34 |





Fig 2(c)


| Bin Number | Spectral Intensity | Bin Number | Spectral Intensity | Bin Number | Spectral Intensity |
|------------|--------------------|------------|--------------------|------------|--------------------|
| 51 | 2 | 183 | 1 | 1025 | 62 |
| 55 | 2 | 195 | 1 | 1026 | 42 |
| 63 | 2 | 197 | 1 | 1027 | 49 |
| 65 | 5 | 198 | 1 | 1032 | 62 |
| 66 | 1 | 212 | 1 | 1034 | 47 |
| 76 | 1 | 213 | 100 | 1041 | 66 |
| 77 | 3 | 214 | 15 | 1046 | 14 |
| 79 | 1 | 215 | 1 | 1047 | 12 |
| 89 | 1 | 228 | 35 | 1048 | 19 |
| 90 | 1 | 229 | 5 | 1053 | 20 |
| 91 | 9 | 580 | 328 | 1056 | 74 |
| 92 | 1 | 590 | 223 | 1059 | 72 |
| 94 | 2 | 664 | 977 | 1060 | 72 |
| 95 | 1 | 677 | 1000 | 1062 | 57 |
| 99 | 4 | 690 | 330 | 1069 | 62 |
| 105 | 1 | 704 | 441 | 1088 | 14 |
| 106 | 2 | 834 | 21 | 1089 | 55 |
| 107 | 4 | 864 | 74 | 1095 | 60 |
| 114 | 1 | 866 | 74 | 1097 | 77 |
| 115 | 1 | 868 | 70 | 1098 | 72 |
| 119 | 19 | 873 | 46 | 1106 | 68 |
| 120 | 2 | 874 | 46 | 1114 | 41 |
| 134 | 1 | 875 | 62 | 1115 | 34 |
| 135 | 4 | 877 | 72 | | |
| 152 | 1 | 883 | 77 | | |
| 165 | 2 | 1009 | 39 | | |
| 169 | 1 | 1011 | 37 | | |
| 181 | 1 | 1019 | 4 | | |

Figure 3 – a hypothetical set of spectrally derived molecular structure descriptors for bisphenol A.

FIG. 4




DISCRIMINANT FUNCTION




COMPONENT 1

00175161.520400

CANONICAL VARIATE 1



Relative Intensity





M/Z

FIG. 6

09106341.020400

COMPONENT FREQUENCY


CANONICAL VARIATE 1




M/Z
FIG. 8(a)

00105311.020400

CANONICAL VARIATE 1



Relative Intensity



M/Z

F1G. 8(6)

09106344.020400

DISCRIMINANT FUNCTION




Fig 9

CANONICAL VARIATE 1





Fig 10


Relative Intensity



M/7.

00105241.020400

DISCRIMINANT FUNCTION



CANONICAL VARIATE 1





Fig 12(a)



CANONICAL VARIATE 1





Fig 12(b)







Fig 13

Fig 14



**COMBINED DECLARATION AND POWER OF ATTORNEY
FOR PATENT APPLICATION**

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled **METHODS FOR PREDICTING THE BIOLOGICAL, CHEMICAL, AND PHYSICAL PROPERTIES OF MOLECULES FROM THEIR SPECTRAL PROPERTIES**, the specification of which

☒ is attached hereto.

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, § 1.56. If this is a continuation-in-part application filed under the conditions specified in 35 U.S.C. § 120 which discloses and claims subject matter in addition to that disclosed in the prior pending application, I further acknowledge the duty to disclose material information as defined in 37 CFR § 1.56 which occurred between the filing date of the prior application and the national or PCT international filing date of the continuation-in-part application.

I hereby claim foreign priority benefits under Title 35, United States Code, § 119(a)-(d) of any foreign application(s) for patent or inventor's certificate or of an PCT International application(s) designating at least one country other than the United States of America listed below and have also identified below any foreign application(s) for patent or inventor's certificate or any PCT International application(s) designating at least one country other than the United States of America filed by me on the same subject matter having a filing date before that of the application(s) on which priority is claimed:

| Prior Foreign Application(s) | | | Priority Claimed | |
|------------------------------|-----------|------------------------|------------------------------|-----------------------------|
| (Number) | (Country) | (Day/Month/Year Filed) | <input type="checkbox"/> Yes | <input type="checkbox"/> No |
| | | | | |

I hereby claim the benefit under Title 35, United States Code, § 119(e) of any United States provisional application(s) listed below:

| Application Number | Filing Date |
|--------------------|-------------|
| | |

I hereby claim the benefit under Title 35, United States Code, § 120 of any United States application(s) or § 365(c) of any PCT International application(s) designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of Title 35, United States Code, § 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, § 1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

| (Application No.) | (Filing Date) | (Status: patented, Pending, abandoned) |
|-------------------|---------------|---|
| | | |

The undersigned hereby authorizes the U.S. attorney or agent named herein to accept and follow instructions from < Name > as to any action to be taken in the Patent and Trademark Office regarding this application without direct communication between the U.S. attorney or agent and the undersigned. In the

event of a change in the persons from whom instructions may be taken, the U.S. attorney or agent named herein will be so notified by the undersigned.

I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application, to file a corresponding international application, and to transact all business in the Patent and Trademark Office connected therewith:

| Name | Reg. No. | Name | Reg. No. |
|-------------------|----------|-------------------|----------|
| James C. Haight | 25,588 | Gloria Richmond | 30,416 |
| Robert Benson | 33,612 | Jack Spiegel | 34,477 |
| Susan S. Rucker | 33,762 | David R. Sadowski | 32,808 |
| John Peter Kim | 38,514 | Steve Ferguson | 38,448 |
| Stephen L. Finley | 36,357 | | |

with an Associate Power of Attorney to the following:

| Name | Reg. No. | Name | Reg. No. |
|------------------------|----------|---------------------|----------|
| Richard J. Polley | 28,107 | Samuel E. George | 44,119 |
| William D. Noonan | 30,878 | Tanya M. Harding | 42,630 |
| Mark L. Becker | 31,325 | Susan Alpert Siegel | 43,121 |
| Donald L. Stephens Jr. | 34,022 | Paula A. DeGrandis | P43,581 |
| Stacey C. Slater | 36,011 | | |

all of the law firm of Klarquist Sparkman Campbell Leigh & Winston, LLP.

Address all telephone calls to William D. Noonan, M.D., telephone number 503/226-7391 and facsimile number 503/228-9446.


Address all correspondence to:
KLARQUIST SPARKMAN CAMPBELL
LEIGH & WHINSTON, LLP
One World Trade Center, Suite 1600
121 SW Salmon Street
Portland, OR 97204-2988

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full Name of First Inventor:

Dwight W. Miller

Inventor's Signature



02/11/00
Date

Residence: Pine Bluff, Arkansas

Citizenship: United States

Post Office Address: 2901 Dan Road
Pine Bluff, AR 71602

Full Name of Second Joint Inventor: Richard Beger

Inventor's Signature

Richard Beger

2/1/00
Date

Residence: White Hall, Arkansas

Citizenship:

Post Office Address: 405 Reynolds #10
White Hall, AR 71602

Full Name of Third Joint Inventor: Jackson O. Lay, Jr.

Inventor's Signature

Jackson O. Lay, Jr.

2/1/00
Date

Residence: Little Rock, Arkansas

Citizenship:

Post Office Address: 6 Pleasant Court
Little Rock, AR 72211

Full Name of Fourth Joint Inventor: Jon G. Wilkes

Inventor's Signature

Jon G. Wilkes

2/1/00
Date

Residence: Little Rock, Arkansas

Citizenship:

Post Office Address: 621 Cumberland #1
Little Rock, AR 72202

Full Name of Fifth Joint Inventor: James P. Freeman

Inventor's Signature

James P. Freeman

2-1-2000
Date

Residence: White Hall, Arkansas

Citizenship:

Post Office Address: 1100 Rock
White Hall, AR 71602